



Factors associated with and mode of delivery in prelabour rupture of membrane at secondary health care

Isharyah Sunarno*, Deviana Soraya Riu, Nasrudin Andi Mappaware

Department of Obstetrics & Gynecology, Faculty of Medicine, Hasanuddin University, Indonesia

ARTICLE INFO

Article history:

Received 28 June 2021

Accepted 30 July 2021

Keywords:

Prelabour rupture of membrane

Mode of delivery

ABSTRACT

Objective: The study aimed to explore factors associated with prelabour rupture of membrane (PROM) and mode of delivery of PROM at secondary health care.

Methods: It was a retrospective case-control study within a year observational period. Data were collected from medical records at St. Khadijah I Mother and Child Hospital Makassar.

Results: There were 4003 samples with 259 cases with PROM. Parity and body mass index (BMI) were factors associated with PROM with $p = 0.000$ and $.032$ consecutively; among them, BMI was more prominent than parity with $OR = 2.392$, and $95\% CI = 1.052-5.442$. Cesarean section is not associated with PROM, $p = .000$.

Conclusion: Primigravidae and obesity were factors associated with PROM, and BMI was more strongly associated with PROM than parity. PROM is not an indication for cesarean section.

© 2021 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Prelabour rupture of membranes (PROM) is a rupture of the amniotic membrane before the labor process. It is a condition frequently found at the end of pregnancy. PROM incidence is about 3% of all pregnancies, with 70% are term pregnancies; it varies about 5–10% of all deliveries. PROM is more frequent in developing countries. The prompt onset of spontaneous labor generally follows ¹⁻⁴ term PROM, and it complicates 8% of pregnancies.^{5,6} Since the exact etiology of PROM is multifactorial and, in some cases yet still unclear, effective treatment for preventing PROM is not certain yet. The impact of PROM, especially on neonatal morbidity and mortality, is noteworthy, so it is pertinent to identify factors associated with PROM in order to anticipate PROM when it happens, to reduce the occurrence of adverse effects of PROM toward maternal and neonatal outcomes, and to promote the health of mothers and neonates. Due to PROM, frequent maternal complications are acute chorioamnionitis, subclinical chorioamnionitis, premature placental separation, and postpartum endometritis. Meanwhile, common complications of PROM to neonates are prematurity (the most common complication), sepsis, asphyxia, and respiratory distress syndrome.^{1,2,4}

Extensive efforts were made to find out factors associated with PROM. A hospital-based study by Addisu et al. (2020) revealed that pathological vaginal discharge, urinary tract infection, vaginal hemorrhage during pregnancy, and history of PROM in a previous pregnancy are factors associated with PROM. In comparison, Endale

et al. (2016) found that PROM duration and maternal residential sites were associated with PROM outcomes. Another study by Eleje et al. (2020) showed that the Chorioquick test could be used as a modality to detect chorioamnionitis in PROM patients.⁷⁻⁹ PROM's high-risk factors categorized as preventable (include maternal smoking, more than three cups of coffee per day in the first trimester, vaginal colonization by: *Chlamydia trachomatis*, *Neisseria gonorrhoeae*, Group B streptococci) and nonpreventable (include the history of PROM, vaginal bleeding in the first and second trimesters, cervical length $<2.5\text{cm}$, positive fibronectin).^{1,2}

PROM remains an important determinant for maternal morbidity and fetal and neonatal morbidity and mortality.⁴ Timely identification of maternal risk factors helps to reduce the incidence and severity of PROM complications. Therefore, it is necessary to study factors associated with PROM and mode of delivery, as such that proper management could be given to mothers with certain risk factors besides given a choice for the appropriate mode of delivery.

Methods

This was a retrospective cross-sectional study within a one-year observational period (January 1, 2019, until December 31, 2019). The population of the study was women in labor. Samples were women who gave birth in St. Khadijah I Mother and Child Hospital. This study was approved by the Health Research Ethical Committee, Faculty of Medicine Hasanuddin University. Data were collected from medical records at St. Khadijah I Mother and Child Hospital implementing consecutive sampling and statistically analyzed using the SPSS program. Bivariate analysis was performed using the Chi-square test, and multivariate analysis was using logistic regression.

Peer-review under responsibility of the scientific committee of the 3rd International Nursing, Health Science Students & Health Care Professionals Conference. Full-text and the content of it is under responsibility of authors of the article.

* Corresponding author.

E-mail addresses: isharyahsunarno@gmail.com, pmc@agri.unhas.ac.id (I. Sunarno).

Table 1
Characteristics of subjects.

Characteristics	PROM		Not PROM		p
	n	%	n	%	
<i>Maternal age</i>					
Low risk	201	5.02	2851	71.22	.594
High risk	58	1.45	893	22.31	
<i>Educational status</i>					
Low level	250	6.25	3621	90.46	.869
High level	9	.22	123	3.07	
<i>Occupational status</i>					
Not working	253	6.32	3604	90.03	.238
Working	6	.15	140	3.50	

Table 2
Factors associated with PROM.

Factors	PROM		Not PROM		p	OR	95% CI
	n	%	n	%			
<i>Parity</i>							
Primigravidae	142	3.55	1173	29.30	.000	.376	.292–.485
Multigravidae	117	2.92	2571	64.23			
<i>Body mass index</i>							
Normal	6	.15	201	5.02	.032	2.392	1.052–5.442
Obese	253	6.32	3543	88.51			

Table 3
Mode of delivery in PROM.

Mode of delivery	PROM		Not PROM		p
	n	%	n	%	
Spontaneous normal delivery	227	5.67	2326	58.11	.000
Cesarean section	32	.80	1418	35.42	

Results

Within the period of the study, there were 4003 samples. Samples were categorized as PROM ($n=259$ subjects; hence the incidence of PROM in 2019 is 6.47% of all deliveries at St. Khadijah I Mother and Child Hospital Makassar) as case group and not PROM ($n=3744$ subjects) as a control group. Inclusion criteria were: PROM subjects were diagnosed by the history of per vaginal fluid leakage and nitrazine test, term pregnancy (>37 weeks of gestational age based on last menstrual period), and complete data on the medical record. Exclusion criteria consist of: patients were referred or dead before delivery management was performed and patients admitted to the hospital on December 31, 2018, and the delivery process occurred on January 1, 2019, and also patients admitted to the hospital on December 31, 2019, and the delivery process occurred on January 1, 2020.

The results of the study were as follows: **Table 1** shows the characteristics of study subjects. Statistical analysis revealed that maternal characteristics based on maternal age, educational status, and occupational status were homogenous.

Table 2 described that parity and Body Mass Index (BMI) were factors associated with PROM, with BMI was the stronger factors related to PROM. The probability of obese pregnant women having PROM was two folds compared to normal pregnant women.

Table 3 revealed that PROM was not an indication for performing a cesarean section.

Discussion

Based on gestational age, PROM was categorized as: term PROM (PROM that occurred at >37 weeks of gestation), preterm PROM

(PROM that happened <37 weeks of gestation), and Previa PROM (PROM before the fetus is viable).¹⁰ As a secondary health care, St. Khadijah I mother and child hospital does not have NICU facilities. Thus preterm cases were referred to health care with more advanced facilities. In this study, all PROM cases were term PROM.

Infection in pregnancy brings about a reduction in membrane tensile strength which eventually caused PROM. Membrane tensile strength reduction was rendered by the collagenolytic effect of metalloproteinases, the effect of bacterial proteases, or repeated stretching by uterine contractions.¹²

In this study, PROM is more frequent in primigravidae than multiparous. This result is in accordance with the study by Ibishi et al. (2015).⁴

In our study, obese mothers were inclined to encounter PROM two folds compared to non-obese mothers. At present, the most common BMI measurement used is the Quetelet index which is calculated:

$$\frac{\text{Weight (kg)}}{\text{Height}^2 (\text{m}^2)}$$

According to World Health Organization (WHO), the classification of obesity was as follows¹¹:

- Underweight: $<18.5 \text{ kg/m}^2$
- Normal: $18.5\text{--}24.9 \text{ kg/m}^2$
- Overweight: $25\text{--}29.9 \text{ kg/m}^2$
- Obese:
 - o Class 1: $30\text{--}34.9 \text{ kg/m}^2$
 - o Class 2: $35\text{--}39.9 \text{ kg/m}^2$
 - o Class 3: $\geq 40 \text{ kg/m}^2$

A study by Fallatah et al. (2019) found that obesity in pregnancy is related to a great amount of maternal and neonatal risk and eventually adverse maternal and neonatal outcomes. The study involved 1037 samples which were classified into 4 classes of obesity. Among several antepartum variables, PROM is one of them.¹² A study by Wendremaire et al. (2011) showed that leptin increase in obese pregnant women has a role in myometrial apoptosis due to leptin-induced inhibition.¹³ The association between prelabour rupture of the membrane with obesity was also found in a study by Hautakangas et al. (2018). Their study that consisted of 598 samples revealed that obesity was an independent risk factor of labor dystocia when BMI was ≥ 30 kg/m² but did not affect neonatal outcome. PROM occurred in 41.9% of cases of severely obese parturients (BMI > 35 kg/m²).¹⁴

We also found that PROM is not an indication for cesarean section. Middleton et al. (2017), in a Cochrane Systematic Review and Metanalysis, proved that there was no clear difference between planned and expectant management for the risk of cesarean section in term pregnancy.⁶ But our result is not in accordance with the study conducted by Ibishi et al. (2015), which revealed an increase in cesarean section delivery due to PROM.⁴ Several factors must be taken into consideration before performing a cesarean section, such as maternal and neonatal morbidity and mortality and subsequent pregnancy and delivery. Clinical possibilities to avoid dystocia in obese pregnant women are limited, so it is worth trying vaginal delivery. For obese parturients, recognizing risk factors is pertinent before deciding to perform a cesarean section.¹⁴

Our study has a strength that is the fairly large study samples from a single center. Thus the homogeneity of the study is good. The data on maternal background and deliveries is reliable, collected from the patients' medical records. The limitation of our study is that some important variables are not included due to incomplete documentation and inappropriate chart keeping. Another limitation is we did not stratified obesity according to WHO classification.

Conclusion

In this study, maternal parity and obesity were factors associated with PROM, with the latter being more strongly related to PROM. PROM is not an indication for cesarean section.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Fernando A, Daftary SN, Bhide AG. Premature rupture of membranes. In: Fernando A, Daftary SN, Bhide AG, editors. *Practical guide to high-risk pregnancy & delivery: a south Asian perspective*. 3rd ed. New Delhi: Elsevier Ltd.; 2008. p. 240–61.
2. Boskabadi H, Zakerihamidi M. Evaluation of maternal risk factors, delivery, and neonatal outcomes of premature rupture of membrane: a systematic review study. *J Pediatr Rev*. 2018;7:77–88.
3. Assefa NE, Berhe H, Girma F, et al. Correction to: risk: factors of premature rupture of membranes in public hospitals at mekele city, Tigray, a case-control study. *BMC Pregnancy Childbirth*. 2018;18:1–7.
4. Ibishi VA, Isjanovska RD. Prelabour rupture of membranes: mode of delivery and outcome. *Maced J Med Sci*. 2015;3:237–40.
5. The American College of Obstetricians and Gynecologists. Prelabor rupture of membranes. *Obstet Gynecol*. 2019;133:168–86.
6. Middleton P, Shepherd E, Flenady V, et al. Planned early birth versus expectant management (waiting) for prelabour rupture of membranes at term (37 weeks or more). *Cochrane Database Syst Rev*. 2017;2017.
7. Addisu D, Melkie A, Biru S. Prevalence of preterm premature rupture of membrane and its associated factors among pregnant women admitted in Debre Tabor General Hospital, North West Ethiopia: institutional-based cross-sectional study. *Obstet Gynecol Int*. 2020;2020:1–7.
8. Endale T, Fentahun N, Gemada D, et al. Maternal and fetal outcomes in term premature rupture of membrane. *World J Emerg Med*. 2016;7:147–52.
9. Eleje GU, Ukah CO, Onyiaorah IV, et al. Diagnostic value of Chorioquick for detecting chorioamnionitis in women with premature rupture of membranes. *Int J Gynecol Obstet*. 2020;149:98–105.
10. Shazly SA, Ahmed IA, Radwan AA, et al. Middle-east OBGYN graduate education (MOGGE) foundation practice guidelines: prelabor rupture of membranes; practice guideline no. 01-O-19. *J Glob Health*. 2020;10:1–17.
11. Uccioli L, Monticone G, Russo F, et al. Autonomic neuropathy and transcutaneous oxymetry in diabetic lower extremities. *Diabetologia*. 1994;37:1051–5.
12. Fallatah AM, Babatin HM, Nassibi KM, et al. Maternal and neonatal outcomes among obese pregnant women in King Abdulaziz University Hospital: a retrospective single-center medical record review. *Med Arch (Sarajevo, Bosnia Herzegovina)*. 2019;73:425–32.
13. Wendremaire M, Bardou M, Peyronel C, et al. Effects of leptin on lipopolysaccharide-induced myometrial apoptosis in an in vitro human model of chorioamnionitis. *Am J Obstet Gynecol*. 2011;205, 363.e1–363.e9.
14. Hautakangas T, Palomäki O, Eidstø K, et al. Impact of obesity and other risk factors on labor dystocia in term primiparous women: a case control study. *BMC Pregnancy Childbirth*. 2018;18:1–8.