



Potency of the *Piper betle* and *Ocimum basilicum* as a natural antibacterial against the acute hepatopancreatic necrosis disease (AHPND)

¹Harlina Harlina, ¹Andi Hamdillah, ²Rosmiati Rosmiati, ¹Muhammad Kasnir, ¹Syahrul Syahrul

¹ Aquaculture Study Program, Faculty of Fisheries and Marine Sciences, Muslim University of Indonesia, Makassar, Indonesia; ² Research Institute for Coastal Aquaculture, Maros, South Sulawesi, Indonesia. Correspondence author: H. Harlina, harlina.harlina@umi.ac.id

Abstract. Herbal extracts are considered as alternative approaches for sustainable aquaculture. The present study aimed to evaluate the antibacterial potential of *Piper betle* and *Ocimum basilicum* against *Vibrio harveyi*, *Vibrio alginolyticus* and *Vibrio parahaemolyticus*. Samples were extracted by the maceration method, then they were examined for their antibacterial activities against *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* by means of paper disc diffusion method. Active extracts were partitioned with different polarity solvent and tested for minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC). The highest diameter of inhibition zone of *P. betle* (dichloromethane extract) and *O. basilicum* (aqueous extract) were 19 and 16 mm, respectively. The MICs and MBCs of the dichloromethane extract of *P. betle* ranged from 0.39 to 0.78 mg mL⁻¹ and from 0.39 to 6.25 mg mL⁻¹, respectively. Meanwhile, the MICs and MBCs of the aqueous extract of *O. basilicum* ranged from 1.56 to 3.12 mg mL⁻¹ and from 3.12 to 12.5 mg mL⁻¹, respectively. The results indicate that the *P. betle* and *O. basilicum* extract seem promising for a sustainable management of the diseases in the aquaculture industry.

Key Words: vibriosis, secondary metabolite, leave extract, extraction, shrimp bacterial pathogen.

Introduction. The fishing industry has experienced a rapid development, especially in the field of aquaculture (Yuan et al 2019). The aquaculture production increased from 3,193,565 tons in 2007 to 16,114,991 tons in 2017 or an increase of more than 5 times (BPS 2020). This is due to the intensification of the aquaculture. However, intensive cultivation can trigger pathogenic infections that attack aquatic organisms. Among the diseases commonly found in aquaculture areas, especially in shrimps, the vibriosis is a real issue (Karthik et al 2014; Sivakumar et al 2014). Vibriosis is known to cause a very high mortality of tiger prawns and vannamei with mortality rates reaching 95% (Nunan et al 2014) and even 100% (De Schryver et al 2014). Shrimp infected by vibriosis are characterized by a slow growth, anorexia, brownish hepatopancreas and red spots on uropods, pleopods and abdomen (Mohamad et al 2019). Several bacteria are known to cause vibriosis: *Vibrio harveyi* (Haenen et al 2014), *Vibrio alginolyticus* (Novriadi 2016) and *Vibrio parahaemolyticus* (Nunan et al 2014) which are also the main cause of Acute Hepatopancreas Necrosis Disease (AHPND). Vibriosis disease can be overcome by giving antibiotics (Harlina et al 2015).

Antibiotics have long been used by farmers in eradicating diseases that infect shrimp. Various brands of commercial antibiotics are circulating in the market such as chloramphenicol, oxytetracycline, kanamycin and various other types, unfortunately commercial antibiotics have a negative impact on shrimp, the environment and consumers (Cabello 2006). Even some export destination countries require that the exported products

they receive do not contain commercial antibiotics (Supartono & Nr 2015). Therefore, other alternatives are needed in treating shrimp infected with vibriosis.

Betel (*Piper betle* L.), known as plant, has been used for their health benefits (Lei et al 2003; Shah et al 2016). *P. betle* leaf can be used as a natural antibacterial plant. The plant have some bioactive compounds such as betal-phenol, chavicol and other phenolic compounds. These compounds are known to have strong potentials as antifungal, antioxidant and antibacteria agent (Lubis et al 2020; Saraswati 2011). The antibacterial activity of betel leaf can inhibit the growth of *Escherichia coli*, *Klebsiella*, *Pasteurella*, *Salmonella* sp., *Staphylococcus aureus* and kill *Candida albicans* (Rasydy et al 2019; Reveny 2011). Betel leaf extract is often used in aquaculture to treat *Aeromonas hydrophila* infected goldfish (*Carrasius auratus*) (Meriyanti 2020; Mulia & Husin 2012), while *Ocimum basilicum* L. (Lamiaceae) is not only known in cooking but also has been used as a medicinal plant for antimicrobial, antioxidant, antinociceptive, antiviral and potential for use in treating cancer others (El-Ashram et al 2017; Khalil 2013; Silva et al 2016). In this study, the antibacterial properties of a herbal extract from the leafs of *P. betle* and *O. basilicum* were evaluated against shrimp pathogenic bacteria (*V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus*).

Material and Method

Sampling. *P. betle* and *O. basilicum* leaves were obtained from Tello Traditional Market, Batua, Makassar City (5°09'00.2"S 119°28'16.8"E). Freshly collected specimen of betel leave and basil were immediately transported to the laboratory. The samples were separated, washed and cleaned thoroughly with tap water and then with distilled water and air-dried in shade and ground into a fine powder at the Integrated Biology Laboratory, Faculty of Fisheries and Marine Sciences, Universitas Muslim Indonesia.

Extraction. *P. betle* and *O. basilicum* leaves were prepared in accordance with the methods described in (Hamdillah et al 2019) with minor modifications. An amount of 500 g of powder for each sample was extracted using the maceration method, with 2.5 L methanol solvent (MeOH) for 24 h, and the process was repeated three times. After filtering through the Whatman No. 1 filter paper, the extracts was concentrated using a rotary evaporator (Heidolph, Germany) (Sachithanandam et al 2021). All the extracts thus prepared were stored for further analysis at 4°C.

Antibacterial assay. Shrimp bacterial pathogen like *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* were used to evaluate the antibacterial activity of extracts from *P. betle* and *O. basilicum*. All isolates were obtained from the Research Institute for Coastal Aquaculture, Maros. The bacteria were cultured and maintained on nutrient agar slants (Oxoid, UK). Bacterial inhibition assay used the diffusion method on double layer agar (Isnansetyo & Kamei 2003). All inoculum of 10^6 cells mL⁻¹ based on standard McFarland 0.5 were transferred into TSB (Oxoid, UK) added with 0.7% agar. Then, the mixtures were poured into TSA (Oxoid, UK). Each sterile paper disc (6 mm diameter, Macherey Nagel, Germany) were impregnated with extracts of a concentration of 50 mg mL⁻¹, placed on the surface of inoculated agar and incubated at 30°C for 24 h. The antibacterial activity was measured based on the diameter of the inhibition zone.

Partition of crude extract. The crude extract of *P. betle* and *O. basilicum* was further partitioned using immiscible solvent: 50 g of active extract was partitioned using solvents with increasing polarity, based onn-hexane, dichloromethane and distilled water, to obtain hexane, dichloromethane and aqueous solutions, respectively. All the organic fractions of the extract (*n*-hexane, dichloromethane and aqueous extracts) were tested for their antibacterial activity as described previously and stored at 4°C prior to use.

Determination of MIC and MBC. The MIC and MBC tests were carried out on the extract with the highest antibacterial activity. Dichloromethane extract of *P. betle* L and aqueous extract of *O. basilicum* were evaluated for MIC and MBC. The MIC and MBC tests were based on a diffusion method on double layer agar (Isnansetyo & Kamei 2003; Isnansetyo & Kamei 2003) and the bacteria tested were *V. alginolyticus*, *V. harveyi*, and *V. parahaemolyticus*. Fractions were tested at different concentrations (50, 25, 12.5, 6.25, 3.13, 1.56, 0.78 and 0.39 mg mL⁻¹), dripped onto a paper disc, then air-dried in an incubator at 30°C before being placed on soft agar media. The diameter of the inhibition zone formed after 24 hours was observed and measured. The lowest concentration of the active extract indicated the zone of inhibition corresponding to the MIC value. On the other hand, the extracts were incubated again at 30°C for 24 h to determine the MBC, as the lowest concentration of drug or extract, which did not produce any bacterial colonies on the agar plate.

Results. In the present study, *P. betle* and *O. basilicum* extracts against the growth of *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* were evaluated. *P. betle* leaves and basil *O. basilicum* leaves were obtained from the Tello Traditional Market, Batua, Makassar City (5°09'00.2"S 119°28'16.8"E). The research on the antibacterial potential of *P. betle* and *O. basilicum* was carried out in several stages. *P. betle* and *O. basilicum* leaves were subjected to an extraction process with methanol solvent and yielding a percentage of 9.9% and 13.93% of crude extract.

Extraction of crude *P. betle* and *O. basilicum*. The antibacterial activity was evaluated by measuring the diameter of the inhibition zone after incubating the plates at 30 °C for 24 h. The antibacterial assay of the bioactive compound of *P. betle* and *O. basilicum* extract showed an efficient suppression of the growth of bacterial disease in shrimp. Leaf extract of *P. betle* and *O. basilicum* had a remarkable antibacterial activity with maximum zones of inhibition about 14 mm against *V. harveyi*, 17 mm against *V. Alginolyticus*, and 16 mm against *V. parahaemolyticus* (Table 1).

Table 1

Antibacterial activities of leaves crude extract from *Piper betle* and *Ocimum basilicum* against different bacterial

Plant crude extract	Diameter of inhibition zone (mm)		
	<i>V. harveyi</i>	<i>V. alginolyticus</i>	<i>V. parahaemolyticus</i>
<i>Piper betle</i>	14	14	16
<i>Ocimum basilicum</i>	14	17	15

Antibacterial activity of the partitioned extract. Crude methanol extract of *P. betle* and *O. basilicum* were fractionated based on the polarity of the compound. The solvents used were *n*-hexane, dichloromethane and distilled water. The *n*-hexane solvent would attract non-polar compounds. Dichloromethane solvent would attract semi-polar compounds and distilled water would attract polar compounds. The antimicrobial activity of the partitioned extract is presented in Table 2. Only the dichloromethane partition extract of *P. betle* showed antibacterial activity. On the other hand, only the aqueous partition extract of *O. basilicum* showed antibacterial activity. Based on the results of the study in table 1. that dichloromethane extract of *P. betle* was able to inhibit the growth of *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* with inhibition zone diameters of 19, 16 and 15 mm, respectively. Meanwhile, aqueous extract and *n*-hexane of *P. betle* did not inhibit the growth of all test bacteria.

Table 2

Antibacterial activity test of extracts of n-hexane, dichloromethane and aqueous extract of *Piper betle* and *Ocimum basilicum* against bacterial pathogens

Plant samples	Extract	Diameter of inhibition zone (mm)		
		<i>V. harveyi</i>	<i>V. alginolyticus</i>	<i>V. parahaemolyticus</i>
<i>Piper betle</i>	Aqueous	-	-	-
	Dichloromethane	19	16	15
	n-Hexane	-	-	-
<i>Ocimum basilicum</i>	Aqueous	16	13	15
	Dichloromethane	-	-	-
	n-Hexane	-	-	-

MIC and MBC. In the current study, the MIC and MBC values were determined from the partitioned extract. The highest antibacterial activity was determined for its minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) (Table 3). The MIC and MBC were evaluated against *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* using a diffusion method on double layer agar (Isnansetyo & Kamei 2003; Isnansetyo & Kamei 2003). Dichloromethane extract of *P. betle* and aqueous extract of *O. basilicum* were evaluated with different concentrations (50, 25, 12.5, 6.25, 3.13, 1.56, 0.78 and 0.39 mg mL⁻¹).

Table 3

Minimum concentration (MIC) test results

Bacteria	MIC (mg mL ⁻¹)		MBC (mg mL ⁻¹)	
	<i>O. basilicum</i> (Aqueous)	<i>P. betle</i> (Dichloromethane)	<i>O. basilicum</i> (Aqueous)	<i>P. betle</i> (Dichloromethane)
<i>V. harveyi</i>	1.56	0.39	3.12	0.39
<i>V. alginolyticus</i>	3.12	0.78	12.5	6.25
<i>V. parahaemolyticus</i>	1.56	0.39	6.25	0.78

Based on Table 3, the MIC value of *P. betle* (0.39 mg mL⁻¹) was lower than the MIC value of *O. basilicum* (1.56 mg mL⁻¹). Likewise, MBC of *P. betle* (0.39 mg mL⁻¹) was lower than MBC of *O. basilicum* (3.12 mg mL⁻¹). The lower the concentration of the extract that can inhibit or kill bacteria, the better the extract. Besides, the MIC and MBC values of the dichloromethane extract of *P. betle* were the same against *V. harveyi*, meaning this bacteria can be inhibited and killed at the same concentration of the plant extract.

Discussion. Shrimp aquaculture is one of biggest industry in the world. Herbal plants, like *P. betle* and *O. basilicum* are known well as natural medicines and may play a role in the development of new phytomedicine drugs to be used for the treatment of diseases. The data summarized showed a broad spectrum of antibacterial activity of *P. betle* and *O. basilicum* against *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus*. The extracts showing a diameter of inhibition zone less than 10 mm were classified as weak antimicrobial (Jang et al 2007) while the extracts with an inhibition zone of more than 15 mm were classified as strong antimicrobials (Al-Madhangi et al 2019). A strong antibacterial activity was shown by *P. betle* against *V. parahaemolyticus* and by *O. basilicum* against *V. alginolyticus* and *V. parahaemolyticus*. *P. betle* and *O. basilicum* had an antifungal and antimicrobial activity against pathogens (gram positive and negative) (Pandey et al 2014; Sarma et al 2018; Voon & Hussin 2014). The ethanolic crude extract of betel leaves is a potential alternative to antibiotics against the *V. alginolyticus* infection (Othman et al 2018). The opportunistic

pathogen *Vibrio* may cause significant losses in crustacean farms (de Souza & Wan 2021). Moreover, AHPND is known to be caused by strains of *Vibrio parahaemolyticus* that contain a plasmid of a unique virulence (Sachithanandam et al 2021). The infection occurs because *V. parahaemolyticus* contains two toxin genes, namely PirA and PirB (Lai et al 2015). These bacteria also trigger the occurrence of white stools or White Feces Disease (WFD) in shrimp (Supono et al 2019). *V. parahaemolyticus* is generally found not only in brackish waters but also in marine waters (Mohamad et al 2019).

Most bioactive compounds isolated from *P. betle*, such as the carvacrol, eugenol, hydroxychavicol and chavibetol (an isomer of the eugenol) are highly volatile and show a poor solubility in the aqueous phase (Kurniasari et al 2021; Othman et al 2018; Zamakshshari et al 2021). They are able to suppress the growth of *Vibrio* sp. (Maharajan & Sajin 2011) and of some fungi like *Candida albicans* (Maharajan & Sajin 2011; Shah et al 2016; Zamakshshari et al 2021) and *Aspergillus niger* (Sarma et al 2018; Zamakshshari et al 2021). Meanwhile, Ahmed et al (2021) reported that *O. basilicum* inhibits the growth of nine known fish pathogens: four gram-positive bacteria (*Enterococcus faecalis*, *Bacillus* sp., *Streptococcus agalactiae* and *S. aureus*) and five Gram-negative bacteria (*Klebsiella pneumoniae*, *A. hydrophila*, and *V. alginolyticus*).

The extract of *P. betle* has phenolic compounds and derivatives that can suppress the growth of *Propionibacterium acne*. Phenolic substances are the most common secondary metabolites with microbial growth inhibitory abilities (Lubis et al 2020). Phenolic compounds suppress the growth of pathogenic microorganisms by inhibiting (bacteriostatic) and killing (bactericidal) the microbes or by denaturing the three-dimensional bacterial proteins (Noventi & Carolia 2016), disrupting the covalent structure of Gram-positive bacteria. It will lead to the destruction of the wall of bacteria cells. Another report explained the activity of the carboxyl group, forming complexes with extracellular and soluble proteins of bacteria, which results in the loss of their infectivity (Lubis et al 2020).

Flavonoid and tannin compounds also contribute to the antimicrobial activity of the *P. betle* leaf extract. The inhibitory effect of tannin is due to the tannic acid. The mechanism proposed was due to its ability to create a change in its potassium concentration, determining the enzymatic reactions' inhibition and the enzyme production's inhibition. The flavonoid mechanism is by disrupting the concentration of potassium in the Gram-positive bacteria which leads to the dysfunction of their cytoplasm membrane (Lubis et al 2020).

On the other hand, the antibacterial activities of *O. basilicum* were presented in Table 2, the best being observed for the aqueous solvent extracts, which inhibited the growth of *V. harveyi*, *V. alginolyticus* and *V. parahaemolyticus* with a maximum diameter of the inhibition zone of 16, 13 and 15 mm, respectively. The extract of *O. basilicum* exhibited a potent inhibitory action on *A. hydrophila*, *A. veronii*, *Pseudomonas fluorescens* and *Streptococcus agalactia* (Dawood et al 2021) and on shrimp pathogens, such as *Pseudomonas aeruginosa*, *S. aureus*, and some *Vibrio* pathogen (Velmurugan et al 2010).

Bioactive compounds contained in *O. basilicum* such as linalool and eugenol are responsible for the antimicrobial activity against three species of gram-positive bacteria (*B. subtilis*, *Clostridium defficile* and *S. aureus*), three species of gram-negative bacteria (*Escherichia coli*, *Klebsiella pneumonia* and *Salmonella typhi*) and three species of fungi (*Aspergillus niger*, *A. flavus* and *Candida albicans*) (Rubab et al 2021; Silva et al 2016). Linalool (an essential oil) is a bioactive compound protecting the cells from the dangerous impact of aflatoxins through two possible mechanisms, firstly by decreasing the DNA binding formation of aflatoxins and secondly by reacting with the ROS increased by aflatoxins (Neveen et al 2017). The antibacterial and antifungal activity of the eugenol in *O. basilicum* can be ascribed to the presence of a free hydroxyl group in the molecule. The hydroxyl group of the eugenol is thought to bind to the proteins, preventing the enzymatic actions. These effect is associated with the antiinflammatory and antioxidant activities. The eugenol has also shown an excellent antifungal activity against dermatophytes, imperfect filamentous fungi and pathogenic yeasts and an efficient activity against a wide range of

gram-negative and gram-positive bacteria (Marchese et al 2017; Snoussi et al 2016). In other reports, the compounds assumed as antimicrobials are the 2,6 octadiene 1,8 diol, phytol, camphor, exo methyl champenilol linalool oxide, cis geraniol, cis carveol (Solikhah et al 2016).

According to Table 3, the herbal extracts showed a different inhibition against the shrimp pathogens. The MIC value of the dichloromethane extract of *P. betle* (0.39 mg mL⁻¹) was better than in the aqueous extract of *O. basilicum* (1.56 mg mL⁻¹). Nevertheless, *O. basilicum* extract is a potential alternative as antibiotic against vibriosis. *P. betle* extract demonstrated the highest effectiveness against *V. alginolyticus* and *V. harveyi*, both with MIC values of 0.39 mg mL⁻¹, whereas against *V. alginolyticus* the MIC values were of 0.78 mg mL⁻¹. *P. betle* can inhibit the growth of *A. hydrophila* with a MIC of 0.37 mg mL⁻¹ (Ahmad et al 2021). Moreover, Nafiqoh et al (2020) reported a MIC of 2 mg mL⁻¹ for the *P. betle* against *A. hydrophila*.

Extracts of *P. betle* and *O. basilicum* can trigger the innate immune response (such as bactericidal activity, serum lysozyme, respiratory burst activity and hematocrit levels) and thus they can reduce the fish and shrimp mortality (El-Ashram et al 2017; Kumar et al 2013; Nafiqoh et al 2020). The right dosage of the methanol extract of *P. betle* is considered optimal for improving the growth performance, enhancing immunity levels and providing antibacterial properties against the selected bacteria in Nile tilapia fish (Mohtar et al 2021). A treatment with the *P. betle* leaf extract increases the survival rate of the catfish to 66.67 to 86.11% compared to only 30.55% for controls (Mulia & Husin 2012). The leave extract of *P. betle* has a significant effect on the survival and on the histopathology of the vaname's hepatopancreas (Annisa 2017). Besides, in vivo analysis showed that the supplemented commercial feed for *P. betle* protects the *Penaeus vannamei* postlarvae against the *V. harveyi* infection after seven days of treatment (Guzman et al 2022) and inhibited the quorum sensing mediated bioluminescence production and biofilm formation in *V. harveyi* (Srinivasan et al 2017). In vivo assessment showed that *O. basilicum* inhibited the growth of five different shrimp pathogens (*A. hydrophila*, *S. aureus*, *P. aeruginosa*, *V. harveyi* and *V. parahaemolyticus*) and improved the survival rate in the *Penaeus monodon* shrimp post-larvae culture (PL 25) (Velmurugan et al 2010).

Conclusions. *P. betle* and *O. basilicum* leaves exhibit bacterial inhibition properties against various pathogens affecting economically important aquaculture species of shrimp. MIC and MBC values of *P. betle* leaves reached 0.39 and 0.39 mg mL⁻¹, respectively. MIC and MBC values of *O. basilicum* leaves reached 1.56 and 3.12 mg mL⁻¹, respectively. The use of *P. betle* and *O. basilicum* extracts seem promising for a sustainable disease management in the aquaculture industry.

Acknowledgements. The authors would like to acknowledge the Directorate of Research and Community Service Fund, Kemenristek-Dikti for their support under e thresearch grant contract: 585/B.03/UMI/IX/2021. The authors would like also to express their gratitude to the Rector of Universitas Muslim Indonesia and also to the chairman of the Research and Human Resource Development and to his staff.

Conflict of interest. The authors declare no conflict of interest.

References

Ahmad S., Daud H. M., Rukayadi Y., Abdelhadi Y. M., Yusoff N. A. H., Kamarudin M. S., 2021 Screening of selected malaysian herbs as the potential broad spectrum antimicrobial agent against bacterial diseases of fish. Malaysian Applied Biology 50(1): 157-168.

- Al-Madhagi W. M., Hashim N. M., Ali N. A. A., Alhadi A. A., Halim S. N. A., Othman R., 2019 Chemical profiling and biological activity of *Peperomia blanda* (Jacq.) Kunth. *PeerJ* 36(1):11-21.
- Annisa N., 2017 The dipping effect of betel leaves extract (*Piper betle*) at different concentration toward clinical symptoms, survival rate, histological and growth of Vannamei (*Litopenaeus vannamei*) infected by *Vibrio harveyi*. *Journal of Aquaculture Management and Technology* 6:1-9.
- Cabello F. C., 2006 Heavy use of prophylactic antibiotics in aquaculture: A growing problem for human and animal health and for the environment. *Environmental Microbiology* 8(7):1137-1144.
- Dawood M. A. O., El Basuini M. F., Zaineldin A. I., Yilmaz S., Hasan Md. T., Ahmadifar E., El Asely A. M., Abdel-Latif H. M. R., Alagawany M., Abu-Elala Nermeen M., Van Doan H., Sewilam H., 2021 Antiparasitic and antibacterial functionality of essential oils: An alternative approach for sustainable aquaculture. *Pathogens* 10(2):1-38.
- De Schryver P., Defoirdt T., Sorgeloos P., 2014 Early mortality syndrome outbreaks: A microbial management issue in shrimp farming. *PLoS Pathogens* 10(4):10-11.
- de Souza Valente C., Wan A. H. L., 2021 *Vibrio* and major commercially important vibriosis diseases in decapod crustaceans. *Journal of Invertebrate Pathology* 181:1-18.
- El-Ashram A., Afifi A., Sakr S., 2017 Effect of basil oil (*Ocimum basilicum*) on nonspecific immune response of Nile-tilapia (*Oreochromis niloticus*). *Egyptian Journal for Aquaculture* 7(2):15-31.
- Guzman J. P. M. D., Yatip P., Soowannayan C., Maningas M. B. B., 2022 *Piper betle* L. leaf extracts inhibit quorum sensing of shrimp pathogen *Vibrio harveyi* and protect *Penaeus vannamei* postlarvae against bacterial infection. *Aquaculture* 547(737452):1-12.
- Haenen O. L. M., Fouz B., Amaro C., Isern M. M., Mikkelsen H., Travers M. A. S., Renault T. S., Wardle R., Hellström A. S., Dalsgaard I., 2014 Virbiosis in aquaculture. *Bulletin of the European Association of Fish Pathologists* 34(4):138-148.
- Hamdillah A., Isnansetyo A., Istiqomah I., Puspita I. D., Handayani D. P., Kaneko T., 2019 Antibacterial activity of coastal plants and marine sponges from Kei Island Indonesia against bacterial fish pathogens. *Pharmacognosy Journal* 11(4):812-817.
- Harlina H., Prajitno A., Suprayitno E., Nursyam H., Rosmiati, 2015 Potential study of kopasanda (*Chromolaena odorata* L.) leaves as antibacterial against *Vibrio harveyi*, disease causative agent of tiger shrimp (*Penaeus monodon* Fabricius) post larvae. *Journal of Aquaculture Research & Development* 6(10):1-5.
- Isnansetyo A., Kamei Y., 2003 MC21-A, a bactericidal antibiotic produced by a new marine bacterium, *Pseudoalteromonas phenolica* sp. nov. O-BC30 T, against methicillin-resistant *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy* 47(2):480-488.
- Jang M. H., Piao X. L., Kim J. M., Kwon S. W., Park J. H., 2007 Antimicrobial activity of clove and rosemary essential oils alone and in combination. *Phytotherapy Research* 21:989-994.
- Karthik R., Jaffar H. A., Muthezhilan R., 2014 Effectiveness of *Lactobacillus* sp. (AMET1506) as probiotic against vibriosis in *Penaeus monodon* and *Litopenaeus vannamei* shrimp aquaculture. *Biosciences Biotechnology Research Asia* 11:297-305.
- Khalil A., 2013 Antimicrobial activity of ethanolic extract of *Ocimum basilicum* leaf from Saudi Arabia. *Biotechnology* 12(1):61-64.
- Kumar S., Raman R. P., Pandey P. K., Mohanty S., Kumar A., Kumar K., 2013 Effect of orally administered azadirachtin on non-specific immune parameters of goldfish *Carassius auratus* (Linn. 1758) and resistance against *Aeromonas hydrophila*. *Fish and Shellfish Immunology* 34(2):564-573.
- Kurniasari S. Y., Tjahjaningsih W., Sianita N., 2021 Antibacterial activity of betel leaf (*Piper betle* L.) leaves extract on *Vibrio harveyi*. *IOP Conference Series: Earth and Environmental Science* 718(1):1-6.

- Lai H. C., Ng T. H., Ando M., Lee C. T., Chen I. T., Chuang J. C., Mavichak R., Chang S. H., Yeh M. D., Chiang Y. A., Takeyama H., Hamaguchi H. O., Lo C. F., Aoki T., Wang H. C., 2015 Pathogenesis of acute hepatopancreatic necrosis disease (AHPND) in shrimp. *Fish and Shellfish Immunology* 47(2):1006-1014.
- Lei D., Chan C.-P., Wang Y.-J., Wang T.-M., Lin B.-R., Huang C.-H., Lee J.-J., Chen H.-M., Jeng J.-H., Chang M.-C., 2003 Antioxidative and antiplatelet effects of aqueous inflorescence *Piper betle* extract. *Journal of Agricultural and Food Chemistry* 51(7): 2083-2088.
- Lubis R. R., Marlisa, Wahyuni D. D., 2020 Antibacterial activity of betle leaf (*Piper betle* L.) extract on inhibiting *Staphylococcus aureus* in conjunctivitis patient. *American journal of clinical and experimental immunology* 9(1):1-5.
- Maharajan A., Sajin C., 2011 Antibacterial activity of herbal plant extracts against the fish pathogen, *Vibrio alginolyticus*. *Planta Activa* 11:154-159.
- Marchese A., Barbieri R., Coppo E., Orhan I. E., Daglia M., Nabavi S. F., Izadi M., Abdollahi M., Nabavi S. M., Ajami M., 2017 Antimicrobial activity of eugenol and essential oils containing eugenol: A mechanistic viewpoint. *Critical Reviews in Microbiology* 43(6):668-689.
- Meriyanti, 2020 [The effectiveness of betel leaf extract (*Piper betle* L.) on the survival of jelawat fish (*Leptobarus hoevenni*) infected with *Aeromonas hydrophila* bacteria]. *Borneo Akuatika* 2(1):20-29. [In Indonesian].
- Mohamad N., Amal M. N. A., Yasin I. S. M., Zamri, Saad M., Nasruddin N. S., Al-saari N., Mino S., Sawabe T., 2019 Vibriosis in cultured marine fishes: a review. *Aquaculture* 512:1-17.
- Mohtar S. H., Rajendran K., Haulisah S., 2021 Immunostimulatory effect of *Piper betle* treatment in nile tilapia (*Oreochromis niloticus*). *Asean Journal of Life Sciences* 1(1):29-40.
- Mulia D. S., Husin A., 2012 [The effectiveness of betel leaf extract in overcoming catfish infected with *Aeromonas hydrophila* bacteria]. *Jurnal Nasional Sainteks* 9(2):22-33. [In Indonesian].
- Nafiqoh N., Sukenda, Zairin M., Alimuddin, Lusiastuti A., Sarter S., Caruso D., Avarre J. C., 2020 Antimicrobial properties against *Aeromonas hydrophila* and immunostimulant effect on *Clarias gariepinus* of *Piper betle*, *Psidium guajava*, and *Tithonia diversifolia* plants. *Aquaculture International* 28(1):1-13.
- Neveen, El-Soud H. A., Mohamed D., El-Kassem L. A., Khalil M., 2017 Chemical composition and antifungal activity of *Syzygium aromaticum* L. essential oil. *Iranian Journal of Medicinal and Aromatic Plants* 33(4):552-561.
- Noventi W., Carolia N., 2016 The potential of green sirih leaf (*Piper betle* L.) for alternative therapy acne vulgaris. *Majority* 5(1):140-145
- Novriadi R., 2016 Vibriosis in aquaculture. *Omni-Akuatika* 12(1):1-12.
- Nunan L., Lightner D., Pantoja C., Gomez-Jimenez S., 2014 Detection of acute hepatopancreatic necrosis disease (AHPND) in Mexico. *Diseases of Aquatic Organisms* 111(1):81-86.
- Othman A. B., Saad M. Z., Haiha N., Yusof N., Abdullah S. Z., 2018 In vitro antimicrobial activity of betel, *Piper betle* leaf extract against *Vibrio alginolyticus* isolated from asian sea bass, *Lates calcarifer*. *Journal of Applied Biology & Biotechnology* 6(04):46-48.
- Pandey A. K., Singh P., Tripathi N. N., 2014 Chemistry and bioactivities of essential oils of some *Ocimum* species: An overview. *Asian Pacific Journal of Tropical Biomedicine* 4(9):682-694.
- Rasydy L. O. A., Supriyanta J., Novita D., 2019 Formulation of 96% ethanol extract of green betel leaves (*Piper betle* L.) in the preparation of anti-acne powder and anti-acne activity test againsts *Staphylococcus aureus*. *Jurnal Farmagazine* 6(2):18-26.
- Reveny J., 2011 Antimicrobial activity of the extract and fraction of red betel leaf (*Piper betle* Linn.). *Jurnal Ilmu Dasar* 12(1):6-12.

- Rubab S., Bahadur S., Hanif U., Durrani A. I., Sadiqa A., Shafique S., Zafar U. Shuaib M., Urooj Z., Nizamani M. M., Iqbal S., 2021 Phytochemical and antimicrobial investigation of methanolic extract/fraction of *Ocimum basilicum* L. Biocatalysis and Agricultural Biotechnology 31:1-10.
- Sachithanandam V., Lalitha P., Parthiban A., Muthukumaran J., Jain M., Misra R., Ramesh R., 2021 A comprehensive in silico and in vitro studies on quinizarin: a promising phytochemical derived from *Rhizophora mucronata* Lam. Journal of Biomolecular Structure and Dynamics 47(2):1-12.
- Saraswati D., 2011 [Effect of betel leaf extract concentration on the inhibitory power of *Escherichia coli*]. Jurnal Healt & Sport 3(2):285-362. [In Indonesian].
- Sarma C., Rasane P., Kaur S., Singh J., Singh J., Gat Y., Dhawan K., 2018 Antioxidant and antimicrobial potential of selected varieties of *Piper betle* L. (Betel leaf). Anais da Academia Brasileira de Ciencias 90(4):3871-3878.
- Shah S. K., Garg G., Jhade D., Patel N., 2016 *Piper betle*: Phytochemical, pharmacological and nutritional value in health management. International Journal of Pharmaceutical Sciences Review and Research 38(2):181-189.
- Silva V. A., Da Sousa J. P., De Luna, Freire Pessôa H., De Freitas A. F. R., Coutinho H. D. M., Alves L. B. N., Lima E. O., 2016 *Ocimum basilicum*: Antibacterial activity and association study with antibiotics against bacteria of clinical importance. Pharmaceutical Biology 54(5):863-867.
- Sivakumar N., Selvakumar G., Varalakshmi P., Ashokkumar B., 2014 *Lactobacillus* sp. A potent probiotic for disease free shrimp aquaculture. International Journal of Recent Scientific Research 5(6):1031-1045.
- Snoussi M., Dehmani A., Noumi E., Flamini G., Papetti A., 2016 Chemical composition and antibiofilm activity of *Petroselinum crispum* and *Ocimum basilicum* essential oils against *Vibrio* spp. strains. Microbial Pathogenesis 90:3-21.
- Solikhah, Kusuma S. B. W., Wijayati N., 2016 Antimicrobial activity of stem and leaf ethanol extract basil (*Ocimum basilicum* L). Indonesian Journal of Chemical Science 5(2):103-107.
- Srinivasan R., Santhakumari S., Ravi A. V., 2017 In vitro antibiofilm efficacy of *Piper betle* against quorum sensing mediated biofilm formation of luminescent *Vibrio harveyi*. Microbial Pathogenesis 110:232-239.
- Supartono W., Nr R., 2015 [Analysis of rejection of Indonesian export products crab (*Portunus pelagicus*) and crab (*Scylla serrata*) in the United States Period 2002 – 2013. Prosiding Seminar Agroindustri dan Lokakarya Nasional FKPT-TPI Program Studi TIP-UTM 1:28-32. [In Indonesian].
- Supono, Wardiyanto, Harpeni E., Khotimah A. H., Ningtyas A., 2019 Identification of *Vibrio* sp. as cause of white feces diseases in white shrimp *Penaeus vannamei* and handling with herbal ingredients in east Lampung regency, Indonesia. AACL Bioflux 12(2):417-425.
- Velmurugan S., Punitha S. M. J., Babu M. M., Selvaraj T., Citarasu T., 2010 Indian herbal medications to replace antibiotics for shrimp *Penaeus monodon* post larvae. Journal of Applied Aquaculture 22(3):230-239.
- Voon W., Hussin M., 2014 Application of betel leaves (*Piper betle* L.) extract for preservation of homemade chili bo. International Food Research Journal 21(6):2399-2403.
- Yuan J., Xiang J., Liu D., Kang H., He T., Kim S., Ding W., 2019 Rapid growth in greenhouse gas emissions from the adoption of industrial-scale aquaculture. Nature Climate Change 9(4):318-322.
- Zamakshshari N., Ahmed I. A., Nasharuddin M. N. A., Mohd Hashim N., Mustafa M. R., Othman R., Noordin M. I., 2021 Effect of extraction procedure on the yield and biological activities of hydroxychavicol from *Piper betle* L. leaves. Journal of Applied Research on Medicinal and Aromatic Plants 24:1-10.
- ***BPS, 2020 [Aquaculture production by province and type of cultivation (tons)]. 2000-2018. [In Indonesian].

Received: 27 October 2021. Accepted: 22 December 2021. Published online: 10 January 2022.

Authors:

Harlina Harlina, Muslim University of Indonesia, Faculty of Fishery and Marine Science, Department of Aquaculture, Jl. Urip Sumoharjo Km. 5, 90231 Makassar, Indonesia, e-mail: harlina.harlina@umi.ac.id

Andi Hamdillah, Muslim University of Indonesia, Faculty of Fishery and Marine Science, Department of Aquaculture, Jl. Urip Sumoharjo Km. 5, 90231 Makassar, Indonesia, e-mail: andi.hamdillah@umi.ac.id

Rosmiati Rosmiati, Research and Development Institute for Coastal Aquaculture, Jl. Makmur Daeng Sitakka 129, 90512 Maros, Indonesia, e-mail: emirosmiati@yahoo.com

Muhammad Kasnir, Muslim University of Indonesia, Faculty of Fishery and Marine Science, Department of Aquaculture, Jl. Urip Sumoharjo Km. 5, 90231 Makassar, Indonesia, e-mail: muhammadkasnir66@gmail.com

Syahrul Syahrul, Muslim University of Indonesia, Faculty of Fishery and Marine Science, Department of Aquaculture, Jl. Urip Sumoharjo Km. 5, 90231 Makassar, Indonesia, e-mail: syahruld@umi.ac.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Harlina H., Hamdillah A., Rosmiati R., Kasnir M., Syahrul S., 2022 Potency of the *Piper betle* and *Ocimum basilicum* as a natural antibacterial against the acute hepatopancreatic necrosis disease (AHPND). AACL Bioflux 15(1):34-43.