



14566-Article Text-28267-1-10-20210330.pdf

Oct 1, 2021

3261 words / 16700 characters

Suharni

Subjective Complaints of Respiratory Disorders Due to Silica Du...

Sources Overview

11%

OVERALL SIMILARITY

1	d.researchbib.com INTERNET	1%
2	arch.uokufa.edu.iq INTERNET	<1%
3	garuda.ristekdikti.go.id INTERNET	<1%
4	"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 CROSSREF	<1%
5	worldwidescience.org INTERNET	<1%
6	ejournal.undip.ac.id INTERNET	<1%
7	media.neliti.com INTERNET	<1%
8	issuu.com INTERNET	<1%
9	L Lukmanulhakim, Andi Achmad Awaludin. "Related factor in lamentation risk of low back pain on inpatient nurse", Jurnal Aisyah : Jur... CROSSREF	<1%
10	elib.stikesmuhgombong.ac.id INTERNET	<1%
11	repository.tudelft.nl INTERNET	<1%
12	www.mysciencework.com INTERNET	<1%
13	Dwiguandi Adi Valan, Agung Kurniawan. "Effect of N95 Masks on Furniture-cutting Workers with Respiratory Symptoms", KnE Life Scie... CROSSREF	<1%
14	journal.fkm.ui.ac.id INTERNET	<1%
15	fkm.unsrat.ac.id INTERNET	<1%
16	garuda.ristekbrin.go.id INTERNET	<1%

17 Merry Sunaryo, Muslika Nourma Rhomadhoni. "ANALISIS KADAR DEBU RESPIRABEL TERHADAP KELUHAN KESEHATAN PADA PEKER... <1%
CROSSREF

18 Muhammad Fadli Ramadhansyah, Nikie Astorina YD, Setiani Onny. "Dust Exposure and Symptoms of Respiratory Disorder on Worker ... <1%
CROSSREF

Excluded search repositories:

- None

Excluded from Similarity Report:

- Bibliography
- Quotes
- Small Matches (less than 8 words).

Excluded sources:

- medicopublication.com, internet, 13%
- www.medicopublication.com, internet, 13%
- www.sciencegate.app, internet, 1%

Original article

Subjective Complaints of Respiratory Disorders Due to Silica Dust Exposure to Workers at PT Bumi Saran Beton Kalla Block

Alfina Baharuddin¹, Suharni A. Fachrin² Astrid Pratiwi Rufaedah Amir³

¹Lecturer in Departement of Environmental Health, Public Health University Moslem Indonesia,

²Asistant Professor in Departement Occupational of, Public Health University Moslem Indonesia,

³Postgraduated Program Magister of Public health University Moslem Indonesia

Abstract

Purpose: The aims to analyze the effect of exposure to silica dust and individual characteristics on subjective complaints of workers'

Methods: The study design used a cross sectional approach. This research was conducted at PT. Bumi Sarana Beton Kalla Block, samples in this study were workers in the ballmill, mixing and packing as many as 27 people using the total sampling method. Data analysis using chi-square test and multiple regression test.

Results: the results showed that the effect of age on subjective complaints of respiratory disorders was $p = 0.904 > 0.05$, the effect of smoking on subjective complaints of respiratory disorders $p = 0.485 > 0.05$, the effect of working tenure on subjective complaints of respiratory disorders $p = 0.728 > 0.05$, the effect of The use of respiratory protective equipment against subjective complaints of respiratory disorders $p = 0.820 > 0.05$, the effect of work history on subjective complaints of respiratory disorders $p = 0.647 > 0.05$, the effect of exposure to silica dust on subjective complaints of respiratory disorders $p = 0.529 > 0.05$.

Conclusion: exposure to silica dust and individual characteristic factors such as age, smoking habits, years of service, use of respiratory protective equipment, and work history are not significant to the subjective complaints of respiratory

Keywords: Silica Dust, Subjective Complaints, Respiratory Disorders.

Introduction

Health problems can arise due to various factors in the work environment, such as: physical, chemical, biological, physiological, and psychological factors¹. The work environment is very easily exposed to chemical factors such as dust, vapors and gases. Dust under certain conditions is a chemical agent that can reduce work comfort, visual disturbances, pulmonary function disorders, and can even cause general poisoning. Continuously inhaled dust can cause lung damage and fibrosis. Dust with a smaller size has a greater potential to cause lung function problems because dust with a size of less than 1μ can enter the alveoli, while dust particles

$<0.1 \mu$ move in and out of the alveoli and do not settle on the alveoli surface²

Data from the ILO suggests that among all occupational diseases, 10% to 30% are lung diseases. It is detected that about 40,000 new cases of pneumoconiosis occur worldwide each year (ILO, 2010). Based on the results of a survey by the Directorate General of PPM & PL in Indonesia, chronic obstructive pulmonary disease ranks first as a contributor to morbidity (35%), followed by bronchial asthma (33%), lung cancer (30%), and 2% others^{3,4,5}.

1602 *Indian Journal of Forensic Medicine & Toxicology, April-June 2021, Vol. 15, No. 2*

One of the health problems caused by the work environment is the disruption of lung function for workers and communities around industrial areas. Various substances can pollute the air such as dust, cement, cotton, asbestos, chemicals, poisonous gases. The effect of dust exposure on labor can cause disturbances, including enjoyment of work, irritation to both the eyes and the respiratory tract and impaired lung function^{1,6,7}. The accumulation of dust in the lungs can reduce the ability of the lungs to inhale and exhale air so that the volume of air inhaled in the lungs is reduced^{4, 8,9}.

One of the products of PT. Bumi Sarana Beton is a lightweight brick, where the materials used in the manufacturing process are silica sand, cement, lime, a little gypsum, water, and aluminum paste as a building material^{10,11}. Where these materials are one of the factors causing the disruption of lung function when continuously exposing workers to certain concentrations, one of which is silica.^{5,12,13} According to the International Agency for Research on Cancer (IARC), silica belongs to group 1 substances that are carcinogenic in humans. Silica is usually found in crystalline form and rarely in an amorphous state⁵. Inhaled silica crystals cause decreased lung function, acute pneumonia, autoimmune disorders, and even lung cancer^{10,14,15}. Silica crystals that settle in the lungs, will oxidize the alveoli walls causing fibrosis. The more silica crystals that settle in the lungs,

the fibrosis that occurs in the alveoli will get worse and cause a disease known as pneumoconiosis silicosis.

The result of the preliminary survey that has been conducted is by interviewing at PT. Bumi Sarana Beton Kalla Block, the total workforce in the three parts of the production process is 27 people. There are several workers who are at risk of direct exposure to dust including complaints from some workers who experience shortness of breath, coughing and sneezing. The workers who are dominant work 12 hours mand it is assumed that the majority of workers breathe dust from the grinding and finishing process. Observations that have been made by researchers are in fact most workers still do not realize the importance of using masks while working.

Materials and Methods

This research is a quantitative study using a cross sectional approach. This research was conducted at PT. Bumi Sarana Beton Kalla Block. The population in this study were workers from the ballmill, mixing, and packing sections. The number of samples in this study were 27 people. Data analysis used chi square test and multiple regression test. The sampling technique used total sampling. The instrument used to measure humidity and air temperature is a thermo / hygrometer, while the wind speed and direction is an anemometer.

Result

Table 1. Distribution of Respondents Based on Individual Characteristics Variables

Variabel	Total	
	n (27)	%
Age		
< 40 years	23	82,5
≥ 40 years	4	14,8
Masa Kerja		
< 3 years	5	18,5
≥ 3 years	22	81,5

Cont... Table 1. Distribution of Respondents Based on Individual Characteristics Variables

Smoking habit		
Smoking	5	18,5
Never smoking	16	59,3
No smoking	6	22,2
Wearing APP		
Always	5	18,5
Seldom	11	40,7
Never wear	11	40,7
Job experience		
dusty	6	22,2
no dusty	15	55,6
no history	6	22,2


Based on table 1 shows that the majority of workers are <40 years old as many as 23 people (82.5%), the most working tenure is ≥ 3 years as many as 22 people (81.5%), the highest result of the respondents' smoking habit is in the category 16 people (59.3%) have ever smoked, the frequency of using APP by workers in the infrequent and non-using categories respectively has the same number, namely 11 people (40.7%), and the highest result of the respondent's work history lies in a not related to dust a number of 15 people (55.6%).

Table 2. Measurement Results of Environmental Physics Parameters


Location	Result assessment		
	Ta (°C)	RH %	Wind velocity (m/s)
Ballmill	31,7	65,1	0,21
Mixing	31,3	69	0,19
Packing	31,6	61,5	0,45

Table 3. Measurement Results of Silica Dust Exposure

Responden	Result assessment
	mg/m ³
Packing1	
Packing2	0,041
Packing3	0,079
Packing4	0,091
Packing5	0,074
Packing6	0,032
Packing7	0,065
Packing8	0,056
Packing9	0,023
Packing10	0,034
Packing11	0,024
Packing12	0,059
Packing13	0,048
Packing14	0,072
Packing15	0,033
Packing16	0,043
Packing17	0,032
Packing18	0,045
Packing19	0,065
Packing20	0,022
Ballmill21	0,052
Ballmill22	0,125
Ballmill23	0,088
Mixing24	0,072
Mixing25	0,013
Mixing26	0,024
Mixing27	0,011
	0,025



(a) Temperature – concentration



(b) humidity - concentration

Figure 1. Effect of Work Environment Physical Parameters on Silica Dust Concentration

In Figure 1a, it can be seen that the effect on dust concentration has a fairly high R^2 value of 0.3795, so it can be said that temperature has a sufficient effect on the concentration of silica dust. In Figure 1b, it can be seen that the effect of humidity on concentration has a very small R^2 value, namely 0.0546, so it can be said that humidity does not have a major effect on the concentration of silica dust.

Table 4 Analysis of the Effect of Variables on Subjective Complaints of Respiratory Disorders

Variable	Subjective Complaints %		p-value
	Yes	No	
Age			
< 40 years	52,2	47,8	0,904
≥ 40 years	75,0	25,0	
Work period			
< 3 years	40,0	60,0	0,728
≥ 3 years	59,1	40,9	
Smoking habit			
Smoking	80,0	20,0	0,485
Never smoking	50,0	50,0	
No smoking	50,0	50,0	
Wearing APP			
Always	40,0	60,0	0,820
Seldom	72,7	27,3	
Never wear	45,5	54,5	
Job experience			
Dusty	66,7	33,3	0,647
No dusty	52,4	47,6	
No history	50,0	50,0	
Silica exposure			
qualify	53,8	46,2	0,529
not qualify	100,0	0,0	

Based on table 4 it can be concluded that the variables age, years of service, smoking habits, use of APP, work history, and exposure to silica dust have no effect on subjective complaints of respiratory disorders in workers.

Discussion

The older a person, the more susceptible to dust exposure. The elasticity of the lungs does not change at the age of 7-39 years, but there is a tendency to decrease at the age of 25 years, and this decrease is more evident after the age of 30 years, so that breathing power

decreases as a result of which the volume of air during breathing will be less. the age variable is significant with respiratory complaints, an increase in age in workers is followed by an increase in the percentage of workers who experience respiratory complaints^{4,8,16,17}.

Smoking habits will accelerate the decline in lung function. The effect of smoke can be greater than the effect of dust which is only one third of the bad effects of smoking. The relationship between smoking and impaired lung function is a dose response relationship. The purpose of the dose response relationship is that the more cigarettes smoked each day and the longer the smoking habit, the risk of developing several respiratory complaints and disease is greater^{9,13,18,19}.

Toxins in cigarettes will accumulate in number in the body, especially in the lungs. The presence of this toxin will inhibit the process of exchanging O₂ gas with CO₂ in the alveoli^{17,20,21}. This condition will worsen as the number of cigarettes smoked increases, alveolar damage is also very likely due to smoking. This will certainly reduce the number of functional alveoli that play a role in the respiration process^{7,17,20}. As a result, there will be a decrease in the function of the lung organs, causing several complaints about breathing such as coughing, phlegm, shortness of breath, and sputum^{8,12,17}.

This is because the workers who actively smoke are only 5 people (18.5%), while the others have ever smoked or have quit smoking, and never smoked at all. In addition, ¹⁸ based on the results of interviews, respondents said that the company prohibited workers from smoking while in the work environment so that the smoking habit did not affect the complaints of respiratory problems by workers during their work^{5,10,22,23}.

Some of the workers also switch to e-cigarettes or what is commonly known as vape, especially younger workers. The use of e-cigarettes does not pose a serious health risk compared to regular cigarettes, which are consumed by burning. vaping does not cause lung problems^{10, 22}. This means that vape is considered safe to use^{5,8,13}. Even consumers who use e-cigarettes on a daily basis are not affected by their lung health. These findings were obtained from physiological, clinical, or inflammatory effects analysis. This study report was analyzed among e-cigarette users aged 23-35 years. It also targets another group of non-smokers of the same

age range^{6,18,13}

But the results of this study contradict previous studies that smoking is a trigger for subjective respiratory complaints. Likewise with the results of the study which states that workers who have a smoking habit have the opportunity to experience complaints and pulmonary function disorders. ¹ Workers who smoke and are in a dusty environment are more likely to experience respiratory problems than workers who are in the same environment but do not smoke.

⁹ The longer a person works, the more he or she has been exposed to the dangers posed by the work environment. Based on the results of research which states that tenure in a company that contains a lot of dust has a high risk of silicosis. Workers who are in an environment with high dust levels for a long time have a high risk of developing obstructive pulmonary disease. In this process, dust pollution in the air is quite high, especially if it is supported by conditions of low room temperature and humidity, which can increase the dust concentration. Workers in this section have worked ≥ 3 years with a length of work ≥ 8 hours per day and based on the results of measurements of respirable dust, the average worker almost exceeds the threshold value, with one respondent who has exceeded the threshold value with complaints of shortness of breath and coughing at the time do its job.

The onset of respiratory problems in industrial workers can greatly depend on the length of exposure and the dose of exposure received. Long exposure to low levels may not immediately indicate respiratory distress. The relationship between exposure and this effect is highly dependent on three things, namely the level of dust in the air, the cumulative exposure dose (the sum of the levels in the air and the duration of exposure) and the residence time or duration of dust in the lungs. Long-term exposure to low levels causes less severe disease than short-term exposure to high levels^{3,7,14}.

Based on the measurement results of respirable dust exposure, it was found that there was only one worker who exceeded the threshold value, namely workers who operated in the ballmill section, the other three people almost exceeded the threshold value. Some of the complaints also came from the ballmill and packing section where the dust came from the manufactured

goods^{14,18}. In a day there are approximately 1000 cubic meters of light bricks produced by the company, the dust produced from the lightweight bricks spreads into the air when workers in the ballmill grind the raw materials, namely silica sand and cement and spill it into containers, as well as in the packing department. who organize and move the production.

Conclusion

exposure to silica dust and individual characteristics such as age, smoking habits, years of service, use of APP, and work history have no effect on subjective complaints of respiratory disorders for workers

Source of Funding - Self-funding

Conflict of Interest- None of the authors has competing interests

Ethical Clearance- taken from ethical committee of Muslim University of Indonesia

References

18. Salawati, Liza. Silikosis. *Jurnal Kedokteran Syiah Kuala* 17 No., 20-26, , 2017. ;2m(65-64
19. Suma'mur. *Higiene Perusahaan Dan Kesehatan Kerja (Hiperkes)*. 2013, Jakarta: Sagung Seto.
20. Novalinda I. Anes, J. M. L Umboh, P. A. T. Kawatu.. Faktor-Faktor Yang Berhubungan Dengan Gangguan Fungsi Paru Pada Pekerja di PT. Tonasa Line Kota Bitung. *Jurnal JIKMU*, 2015, Vol. 5, No. 3.
21. Windari, Diah, Faktor-Faktor Yang Berhubungan Dengan Gangguan Fungsi Paru Pada Pekerja Bagian *Refinery* Di Pt.Antam Tbk Ubpn Sultra. 2016, ;5(2) 100-125
22. Putu E, Bagus Ida, Putu Andrika. Hubungan antara Paparan Debu Silika dengan Transforming Growth Factor- β 1 Serum pada Pekerja Industri Pengolahan Batu. *Jurnal Penyakit Dalam Indonesia*. 2019, Vol. 6, No. 2.
23. Departemen Kesehatan Republik Indonesia. *Pengawasan kualitas Kesehatan Lingkungan dan Pemukiman*. Dirjen P2M & PLP : 1994, Jakarta.
24. Regia, Rinda Andhita. Katharina, Oginawati. Potensi Bahaya Debu Silika Terhadap Kesehatan Pandai Besi Desa Mekarmaju Kabupaten Bandung. *Jurnal Dampak Teknik Lingkungan UNAND* 2017, 14(2): 73-80.
25. Loscalzo, J. *Pulmonologi dan Penyakit Kritis*. 2nd Edition. Jakarta: GC.
26. Rismandha, Rifqi, dkk. Analisis Pengaruh Faktor-Faktor Risiko Gangguan Fungsi Paru pada Pekerja Area Produksi Industri Kayu. *Proceeding 1st Conference on Safety Engineering and Its Application*.2018, ISSN No. 2581 – 1770 18(4) 92-98
27. Khumaidah, Analisis Faktor-Faktor Yang Berhubungan Dengan Gangguan Fungsi Paru Pada Pekerja Mebel PT Kota Jati Furnindo Desa Suwawal Kecamatan Mlonggo Kabupaten Jepara. Tesis. Semarang : Magister Kesehatan Lingkungan, Program Pascasarjana. 2008. Universitas Diponegoro ;5(2):60-65
28. Anggar, Devi, Corie I. Kualitas Fisik Dan Kimia Udara, Karakteristik Pekerja, Serta Keluhan Pernapasan Pada Pekerja Percetakan Di Surabaya. *Jurnal Kesehatan Lingkungan* Vol. 8, No. 2 Juli 2015: 195-205.
29. Martinus G, Yunus F, Antariksa B. Faal Paru pada Polisi Lalu Lintas Jakarta Pusat dan Faktor – Faktor yang Mempengaruhi. *J Respir Indo*. 2015;35(2):97-106.
30. Nugraheni, dkk. Analisis faktor resiko kadar debu organik di udara terhadap gangguan fungsi paru pada pekerja industry penggilingan padi di Demak. *Jurnal Kesehatan Lingkungan Indonesia* 2014, Vol.3 No.2.
31. Aryaningsih, Siti, dkk. Faktor Risiko Keluhan Subyektif Gangguan Pernapasan Pada Petugas Penyapu Jalan. *Jurnal Ilmiah Permas: Jurnal Ilmiah STIKES Kendari* Volume 10 No 1, Hal 109 – 114.
32. Jacobsen, G. dkk. Longitudinal Lung Function Decline and Wood Dust Exposure in the Furniture Industry. *European Respiratory Journal*.. 2015.
33. Utama, P. A. Hubungan Antara Masa Kerja dan Penggunaan Alat Pelindung Diri Dengan Kapasitas Vital Paru Pada Pekerja Unit Spinning. *Unnes Journal of Public Health* 2015. ;5(2):66-70
34. Mengkidi D, Nurjazuli, Sulistiyani .Gangguan Fungsi Paru dan Faktor-faktor yang Mempengaruhinya pada Karyawan PT. Semen Tonasa Pangkep Sulawesi Selatan. *Jurnal Kesehatan Lingkungan Indonesia*. 2016;5(2):59-64
35. Putu E, Bagus Ida, Putu Andrika. Hubungan antara Paparan Debu Silika dengan Transforming Growth

1608 *Indian Journal of Forensic Medicine & Toxicology, April-June 2021, Vol. 15, No. 2*

- Factor-β1 Serum pada Pekerja Industri Pengolahan Batu. *Jurnal Penyakit Dalam Indonesia*. 2019, Vol. 6, No. 2. ;5(2):41-45
36. Rismandha, Rifqi, dkk, Analisis Pengaruh Faktor-Faktor Risiko Gangguan Fungsi Paru pada Pekerja Area Produksi Industri Kayu. *Proceeding 1st Conference on Safety Engineering and Its Application*. 2018, ISSN No. 2581 – 1770. 10n(2):175–180
37. Eka, Yudha. Analisis Faktor Fisik Lingkungan Dan Karakteristik Pekerja Dengan Keluhan Pernapasan Pada Pekerja Di Industri Panci Aluminium. *Jurnal Kesehatan Lingkungan*, 2018, Vol.10, No.4. ;3(2):20-26
38. Sahli, Z., Pratiwi, R. L. Hubungan Perilaku Penggunaan Masker Dengan Gangguan Fungsi Paru Pada Pekerja Mebel Di Kelurahan Harapan Jaya Bandar Lampung. Bandar Lampung: Sekolah Tinggi Ilmu Kesehatan. 2017, vol 2 no 3. ;5(2):60–65
39. Iriyanti A, Nurjazuli, Suwondo A. Hubungan Kadar Debu Terhirup (*Respirable*) dengan Kapasitas Vital Paksa Paru pada Pekerja Mebel Kayu di Kota Jayapura. *J Kesehatan Lingkungan Indonesia*. 2012;11(2):182–6.
40. Kesuma A, Rachmawati S, Seviana R. Faktor-Faktor Yang Berhubungan Dengan Gejala Gangguan Sistem Pernapasan Akibat Paparan Debu Silika (SiO₂) Pada Area Hand Moulding I, Hand Moulding II, Hand Moulding III, Felling Dan Melting Pekerja Pabrik 1 Pengecoran PT. Barata Indonesia (Persero). 2018,. *Jurnal Kesehatan*. ISSN 2620-7761. Vol. 11. No. 1. 11(2):182–6