Bukti Korespondensi

Nama Penulis : Ernaningsih

Judul Artikel : Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi

Nama Jurnal : Egyptian Journal of Aquatic Biology & Fisheries

Correspondent Process

Tahapan	Keterangan Kegiatan	Tanggal Kegiatan
1	Editorial system registeration	12 Maret 2024
2	Submitted Manuscript to the journal “Egyptian Journal of Aquatic Biology and Fisheries”	13 Maret 2024
3	Manuscript Needs Revision	13 April 2024
4	Manucript Attached	13 April 2024
5	First revision	15 April 2024
6	Second revision submitted	16 April 2024
7	Paper accepted	27 April 2024
8	Paper published	23 Mei 2024

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Editorial system registeration

1 message

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg> Tue, Mar 12, 2024 at 10:40 PM
Reply-To: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
To: ernaningsih.aras@umi.ac.id

Dear Ernaningsih Ernaningsih,

Welcome to **Egyptian Journal of Aquatic Biology and Fisheries** website.

Thank you for your registration in the Editorial System Online Subbmission and peer review tracking system.
Below please find your username and confidential password, which you need to access the system at:

https://ejabf.journals.ekb.eg/

Your username is: **#ErnaNingsih23**

Your password is: **t64wsrkxu1**

Please save this information in a safe place.

Once you login , you can change your password on your profile.

Editorial Office

Egyptian Journal of Aquatic Biology and Fisheries

https://ejabf.journals.ekb.eg/

Manuscript ID	EJABF-2403-4089 (R1)
Manuscript Title	Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi
Manuscript Type	Original Article
Main Subjects	Fish Biology
Abstract	A total of 1315 individuals were collected from Bawasalo Waters during the period March – August 2023 From the catch of fishermen with bubu naga fishing gear. This study aims to analyze population parameters and exploitation rate of Blue swimming crab (Portunus pelagicus) who was caught with a bubu naga in the waters of Bawasalo, Pangkep District, South Sulawesi.. The observed parameters include the size distribution of the carapace width , carapace width-weights relationship, Age group, Growth rate , total mortality rate, natural rate, fishing rate and exploitation ratio analyzed with the ELEFAN II program. Size carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as W = 0.1038L2.7334 (negative allometric), The Von Bertalanffy growth parameters were estimated to be K = 0,41 yr-1, L maks = 15.10 cm, t0 = -0,723 yr, value total mortality (Z), natural mortality (M) and fishing mortality (F) were calculated as 2,70, 1.22, 1.48 yr-1 respectively. Exploitation rate (E = 0.55) indicates that the stock of P. pelagicus was slightly higher than that optimally exploited stock
Keywords	Portunus pelagicus, growth, mortalities, exploitation ratio, Bawasalo Waters
Suggested Reviewers	Tahya, Akbar Marzuki
Comments	dear I hope my journal can be accepted and get a response quickly. and can be published

Authors

#	Name	Email Address	Degree	Position	Phone	Country	Affiliation
1	Ernaningsih, Ernaningsih*	ernaningsih.aras@umi.ac.id	PhD	Associate Professor	081355253723	Indonesia	Faculty of Fisheries and Marine Sciences. Indonesian Muslim University
2	DANIAL, DANIAL	danial.danial@umi.ac.id	PhD	Associate Professor	08124160315	Indonesia	Marine Science, Fisheries and Marine Science. Universitas Muslim Indonesia Makassar
3	Syarul, Syahrul	syahruld@umi.ac.id	PhD	Instructor	+62 821-8818-4454	Indonesia	Aquaculture Study Program, Faculty of Fisheries and Marine Science, Universitas Muslim Indonesia, Indonesia

4	Moh. Tauhid Umar, Moh. Tauhid Umar	attauhid@unhas.ac.id	PhD	Associate Professor	08124169514	Indonesia	Faculty of Marine Science and Fisheries, Hasanuddin University Makassar
5	syahran, Nursyahran	nursyahran00@gmail.com	PhD	Associate Professor	085255233596	Indonesia	Institute of Maritime Technology and Business Balikdiwa

Submit Date	2024-03-13 11:09:35
Current Status	Manuscript Needs Revision (Minor Revision)
Modify Date	2024-05-08 08:39:01

Related Files

#	File Type	File Name	Size	File Description	Upload Date
Files Sent by Authors					
1	Title Page	1. Title Page.docx	30.45 KB		2024-03-13
2	Manuscript Main File	2. Manuscript Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regen.docx	60.24 KB		2024-03-13
3	Figure	3. figure.docx	1.07 MB		2024-03-13
4	Table	4. Table.docx	67.96 KB		2024-03-13

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Acknowledgement of Submission (Manuscript #EJABF-2403-4089)

1 message

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg> Wed, Mar 13, 2024 at 4:09 PM
Reply-To: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
To: ernaningsih.aras@umi.ac.id
Cc: ejabf2017@gmail.com

Manuscript ID: EJABF-2403-4089

Manuscript Title: **Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi**

Authors: Ernaningsih Ernaningsih, DANIAL DANIAL, Syahrul Syarul, Moh. Tauhid Umar Moh. Tauhid Umar, Nursyahran syahran

Dear **Dr. Ernaningsih Ernaningsih**

I wish to acknowledge receiving the of the above mentioned manuscript.

It should be noted that the manuscript will be reviewed for possible publication in the Scientific Journals Management System.

Please be sure that the submitted manuscript has not been published previously and will not be submitted elsewhere prior to our decision.

Our editorial decision will be brought to your attention once the paper has been reviewed due the referees consideration.

I wish to take this opportunity to thank you for sharing your work with us.

Truly yours,

Executive managing Editor of **Egyptian Journal of Aquatic Biology and Fisheries**

- Editor-in-Chief Comments
- ✓ Select Manuscript Type
- ✓ Enter Manuscript Title
- ✓ Add/Remove Authors
- ✓ Enter Abstract
- ✓ Enter Keywords
- ✓ Select Related Subjects
- ✓ Additional Comments
- Attach Files
- ✓ Cover Letter & Checklist
- Finish Submission

Editor-in-Chief Comment for Author

Reviewers Recommendation

Reviewer 1	EJABF-2403-4089	Minor Revision

Reviewer Comment For Author:

Please follow the instruction for revision in manuscript comment.

Manuscript Evaluation Form

File Sent by Reviewer:
EJABF-2403-4089-3-1.docx
EJABF-2403-4089-2-2.docx

Original Manuscript Status: Manuscript Needs Revision (Minor Revision)

Revision Due Date: 2024-04-28

Editor-in-Chief Comment

Dear Author,
Please use the attached format template of the Journal (that is present in the Guide for Authors section) after correcting your manuscript and stick to all instructions to save time on publishing, and send it to mtkhalil52@hotmail.com, and put the number of the article.
please save it in word 97/2003
Regards,
EJABF

Next

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Manuscript Needs Revision (Manuscript #EJABF-2403-4089 (R1))

1 message

Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg> Sat, Apr 13, 2024 at 5:20 PM
Reply-To: Egyptian Journal of Aquatic Biology and Fisheries <jssub@ekb.eg>
To: ernaningsih.aras@umi.ac.id

Manuscript ID: EJABF-2403-4089

Manuscript Title: **Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi**

Authors: Ernaningsih Ernaningsih, DANIAL DANIAL, Syahrul Syarul, Moh. Tauhid Umar Moh. Tauhid Umar, Nursyahran syahran

Dear **Dr. Ernaningsih Ernaningsih**

Evaluation process of the above mentioned manuscript has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s) comments and revise your manuscript within the period of defined time.

Because we are trying to facilitate timely publication of manuscripts submitted to journal, your revised manuscript should be uploaded as soon as possible. If it is not possible for you to submit your revision in a reasonable amount of time, we may have to consider your paper as a new submission.

Once again, thank you for submitting your manuscript to this journal and I look forward to receiving your revision.

Truly yours,

Editorial Office of **Egyptian Journal of Aquatic Biology and Fisheries**

Dear Author,

Please use the attached format template of the Journal (that is present in the Guide for Authors section) after correcting your manuscript and stick to all instructions to save time on publishing, and send it to mtkhalil52@hotmail.com, and put the number of the article.

please save it in word 97/2003

Regards,

EJABF

Reviewers Recommendation:

Reviewer 1:

File Sent by Reviewer:
https://ejabf.journals.ekb.eg/jufile?__file=JzrEmIVJYLk8GZyrb1tg2J3eprxxoRDY7btF_cCzGOJGe789e7CDfiWnyj.G3UJV.jt2C9rpd6QNr5HjoxZhPL3sg9Z_STz_Etqi2jaGUN8KmPScHXDU5tnLZInJvs.Q9SYAL4M6YtSC0ie9VlIIFG9QFcW4yQFxr30YyIIzfh.Zg6I.XVpeIlS2FZHwsqOH
https://ejabf.journals.ekb.eg/jufile?__file=gtr05nkG2kErt1Cx3aYIhndTsKMwu3gUZTwZqpw1fpKSS4eDoagrHV9W4rXLMi8l4Au2h5oW8v6Ody3tZl7NNZr_lcP8Ot2_P_LTZFV2EdK33FY9Dthcbz4ylTn.bLGRwjhygjMbF8t6KeMFhLXP5fq_GKjm4rEH4ViwmvB3ADZc.OPwKmAum1puOsPxcQWy
Reviewer Comment For Author:

Please follow the instruction for revision in manuscript comment.

Manuscript Evaluation Form:
https://ejabf.journals.ekb.eg/author?__au=X33PmRXrSLim9ahvskYI6fxPI5saXo4AWOvfATl6I6.6b5GgrQzW1gLmsjjriYEm

Population parameters and exploitation ratio of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi

ARTICLE INFO

Article History:
Received:
Accepted:
Online:

Keywords:
Portunus pelagicus, growth, mortalities, exploitation ratio, Bawasalo Waters

ABSTRACT

A total of 1315 individuals were collected from Bawasalo Waters during the period March – August 2023 From the catch of fishermen with bubu naga fishing gear. This study aims to analyze population parameters and exploitation rate of Blue swimming crab (*Portunus pelagicus*) who was caught with a bubu naga in the waters of Bawasalo, Pangkep District, South Sulawesi.. The observed parameters include the size distribution of the carapace width , carapace width-weights relationship, Age group, Growth rate , total mortality rate, natural rate, fishing rate and exploitation ratio analyzed with the ELEFAN II program. Size carapace width 6.70 ~~cm~~ – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), The Von Bertalanffy growth parameters were estimated to be K = 0,41 yr^{-1}, L maks = 15.10 cm, t0 = -0,723 yr, value total mortality (Z), natural mortality (M) and fishing mortality (F) were calculated as 2,70, 1.22, 1.48 yr^{-1} respectively. Exploitation rate (E = 0.55) indicates that the stock of *P. pelagicus* was slightly higher than that optimally exploited stock

INTRODUCTION

Blue swimming crab is widely distributed throughout Indonesia (**Kusuma dan Dewi, 2017**). In several regions in Indonesia, blue swimming crab is used as one of the people's fishery businesses including the Thousand Islands (**Agus *et al*., 2016**), Lasongko Bay (**Hamid *et al*., 2015**;) and Pangkep Regency **(Ihsan *et al*., 2014**; **Nurdin *et al*., 2016**)

Pangkep Regency located in South Sulawesi has waters with considerable blue swimming crab (*P. pelagicus*) potential and became one of the important commodities. In Pangkep Regency itself, one of the blue swimming crab producing centers is on Salemo Island (**Nurdin *et al*., 2016b**). Total exports of blue swimming crab in 2020 reached 27,616,332 tons, with export value reaching USD 367,519,713 (**KKP, 2021**) is fourth in the total export value of Indonesian fishery products after shrimp, tuna and squid/cuttlefish/octopus

The high export value indicates high production activities and exploitation of blue swimming crab, supported by open access fishing activities. Blue swimming crab (*P. pelagicus*) is currently under considerable pressure as the volume and size of the catch decreases (**Wiyono, 2015**). High fishing pressure from blue swimming crab (*P. pelagicus*) in some areas in Indonesia has led to reduced stocks (**Tirtadanu, 2019**). The catch and abundance of blue swimming crab on Salemo Island, Pangkep Regency has decreased and affected the income of fishermen (**Nurdin *et al*, 2015**). Fisheries resources have seriously declined due to overfishing, illegal fishing activities and pollution **(Lin *et al*., 2019) ; Mehanna *et al*., 2018**) **; Asiedu *et al*., 2023**)

Commented [AMT'S1]: Dynamics

Commented [AMT'S2]: Many references use "mm", i suggest for your consistency and can be easily to compare.

Commented [AMT'S3]: Many references use "mm", i suggest for your consistency and can be easily to compare.

~~Biological information and~~ population dynamics are very important for stock estimation (**Then *et al*, 2018**). Information about biological aspects is a major factor for management purposes (**Sujatha *et al*, 2015**). For management purposes, information on fisheries biology and population structure is required (**El-Kasheif, 2021**). The sustainability of the blue swimming crab requires population biology information and estimating fishery stocks as a scientific basis study for the availability of sustainable resources **(I. Maiyza *et al*., 2020; Froese *et al*, 2011**). Research related to blue swimming crab in Pangkep Regency has been made regarding the arrest (**Ihsan, 2018).** Bawasalo waters, is one of the blue swimming crab fishing areas in Pangkep Regency, However a study of population parameters and exploitation ratio of blue swimming crab (*P. pelagicus*) those caught with bubu naga in Bawasalo waters have not been found. Based on this, it is important to conduct research related to population parameters and exploitation rates which include the distribution of carapace width, carapace width- weight relationship, age group, growth rate, mortality rate dan rate of exploitation as one of the important studies in the sustainability of blue swimming crab resources.

MATERIALS AND METHODS

Total of 1159 blue swimming crabs (*P. pelagicus*) randomly drawn from March 2023 to August 2023 is the catch of bubu naga originating from Bawasalo Waters, Segeri District, Pangkep Regency (**Fig. 1**). Each sample of blue swimming crab measured carapace width (CW) in cm (The width of the blue swimming crab is measured from the longest lateral spines on the sides of its body) and weight (W) in g.

Figure 1. Map of research location *P. pelagicus* in Bawasalo waters

Length frequency data was analyzed by Bhattacharya's method (**Bhattacharya, 1967**) that was involved in the FiSAT II software package (**Gayanilo, 2005**).

The total length and body weight of fish were used for the length-weight relationship. The length-weight relationship was analyzed using the equation below (**Ricker, 1975**):

$$W = aL^b$$

Where, W = Total body weight of the fish (g); L = Total carapace width (cm); "a" = Regression constant, and "b" = Koefisien regresi; Value "b" provide information about the type of fish growth. The degree of association between length-weight variables is calculated by the correlation coefficient (R2), and the degree of its statistical significance is assessed following the method indicated by **Santos *et al*. (2002)**, as follows:

$$\text{Log } W = \text{Log } a + b \text{ Log } TL$$

If b = 3, The growth is isometric, $b \neq 3$, The growth is allometric (allometric negative if $b < 3$ and allometric positive if $b > 3$).

Commented [AMT'S4]: P

Commented [AMT'S5]: Basic scientific studies

Commented [AMT'S6]: why do you use the lateral spine?, instead this part will be very fragile and cause biased data.

It's better to use a body curve that is located near the longest spine.

Commented [AMT'S7R6]: we suggest for easily to compare, if you use mm.

Commented [AMT'S8]: In your map please provide "Map Legend" and "Map Source".

Growth parameters follow the growth function of Von Bertalanffy (VBGF) including growth rate (K) and asymptotic length (L∞) with equations.

$$L_t = L_\infty\left(1 - e^{-K(t - t_0)}\right)$$

Where, Lt is the average length at time (or age) t, L maks; is the asymptotic mean length; K is the coefficient of growth rate (per year), t; represents time or age. The theoretical age at zero length (t0) follows the equation shown by **Pauly (1979)**, as follows:

Log 10 (-t0) = -0.3922 – 0.2752 log 10 L∞ - 1.038log 10 K

Estimation of total mortality (Z) was carried out by the length converted catch curve method **(Pauly, 1983)**. Natural mortality rate **(M) = Log M = -0.0066 – 0.279 log L∞ + 0.6543 log K + 0.4634 log T (Pauly, 1980),** with a flat temperature of 30^0 C. Fishing mortality rate (F)= Z-M. Exploitation rate using equations **Gulland (1971)**, **E = F/Z**

Long was first caught allegedly based on probabilitas fishing cumulative against the middle of the width class. From the resulting curve, Long first caught (Lc50) which is defined as the average total length where 50% of the fish are caught, taken according to the cumulative probability at 50%. This procedure follows the method by **Beverton dan Holt (1956**) dan **Sparre dan Venema (1998)**, as follows :

Lc50 = [TL – K(TL∞ – TL‟)]/ Z

Where , Lc50 is the first length, and T times caught where L 'is the average length of fish in the catch sample. Moreover, length at 75 and 95% fishing correlates with cumulative probability at 75 and 95%, respectively, as shown by **Pauly (1983).**

The result per recruit (Y / R) is determined according to the formula developed by **Beverton dan Holt (1957**), dan hasil relatif per rekrut (Y / R0) dihitung menurut **Pauly dan Soriano (1986).**

Data analysis

The FiSAT II tool was used to assess the population parameters, such asymptotic length (L∞), growth rate (K), natural mortality rate (M), fishing mortality rate (F), exploitation ratio (E), and relative yield per recruit indicators of samples from the assessed fish species encountered during the study period (**Gayanilo *et al*. 2005)**

RESULTS

Commented [AMT'S9]: Please provide "Gonad Maturation Stages" for your result.

Carapace width distribution

Carapace width distribution of blue swimming crab carapace (*P. pelagicus*) during the study from March 2023-August 2023, Who was caught with the bubu naga in Bawasalo waters Pangkep Regency to 1315 individuals on the group width carapace 6,70 ~~cm~~ – 13,80 cm, with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) (**Fig. 2**).

Fig (2). Carapace width distribution of blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

Carapace Width-weight relationship

The carapace width and weight relationship for males, females and combined sexes of *P. pelagicus* was described by equation as : $W= 0.1052L^{2.7245}$ (R2 = 7478), **(Fig 3**)

Fig (3). Width-weight Relationship of blue swimming crab (*P. pelagicus*) in Bawasalo waters

Age Group

Cohort analysis with Battacahrya's method *P. Pelagicus Consists of 3 age groups* is shown in **Fig (4).** Three cohort group were identified at length 8.66 sd 0.67, 10.27 sd 0.93, 12.16 sd 0.21 cm

Fig (4). Cohort groups by Bhattacharya's methode in Bawasalo Waters

Growth parameters

Estimated values of growth parameters were analyzed using the FISAT II program, The results of the analysis of the growth parameters estimation with the formula von Bertalannffy of blue swimming crab *P. pelagicus* including maximum length ($L\infty$), growth coefficient (K) and age at birth (t0) that is $L\infty = 15.1$ cm, K = 0.41 per month and t0 = -0.48 **(Fig 5).**

Fig. (5). Growth curve of Von Bertalanffy blue swimming crab (*P. pelagicus*) in Bawasalo waters

Moralities and exploitation ratio

Commented [AMT'S10]: Mortalities

Mortality analysis based length-converted catch curve (**Fig. 6**) obtained total mortality rate, natural mortality, fishing mortality dan exploitation ratio blue swimming crab (*P. pelagicus*).

Fig. (6). Growth curve of Von Bertalanffy blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

The total mortality (Z) was 2.70 yr^{-1}, natural mortality (M) and fishing mortality (F) was 1.22 yr^{-1} and 1.48 yr^{-1} and the exploitation ratio was 0.55 (**Table 1**).

Table 1. Value of population paramters blue swimming crab (*P. pelagicus*) in Bawasalo waters

Length at first capture

The length first capture (Lc50) is 10.92 cm. Length of catch probability 25% (Lc25) and 75% (Lc75) as 10.53 cm dan 11.22 cm **(Fig. 7).**

Fig. (7). Probability of capture

Yield per recruit (Y/R), and relative yield per recruit (Y'/R)

The yield per recruit (Y/R) for P. pelagicus was 20.6 gr pada current level mortality (F cur) 4.8 yr^{-1} (**Fig 8**). Relative yield per Y'/R recruit for different exploitation ratio (**Fig 9)**. From this curve, the maximum Y'/R is found at the exploitation ratio (Emax) = 0,50. The exploitation ratio E = 0.3.. would give the maximum Y/R' results in the protection of about 50% of the stock biomass

Fig. (8). Yield per recruit, biomass per recruit of blue swimming crab (*P. pelagicus*) in Bawasalo waters

Fig. (9). Relative yield per recruit and relative biomass per recruit in relation to different exploitation ratio of *P.pelagicus* in Bawasalo Waters

DISCUSSION

Carapace width distribution of blue swimming crab (*P. pelagicus*) during the study that is 6,70 cm – 13,80 cm, where the size of the carapace width < 10 cm as 66,69% and carapace width >10 cm as 33,31%**.** Research results **Ihsan (2018)** in Pangkep's waters finding the dominant class interval males blue swimming crab on size 112-123 mm with a frequency of 265 heads (31,77%) and blue swimming crab females at intervals of classes 100-111 mm with a frequency of 223 heads (26,74%). Research results **Abdul Hamid (2019**) in Lasongko Bay shows carapace width on male blue swimming crab ranges from 4.1 cm -12.4 cm. Size of *P. pelagicus* obtained in Kwandang waters ranges from 7.0 – 17.5 cm (**Tirtadanu *et al*., 2019**). Blue swimming crab catches obtained in Gresik and Lamongan dominated by the size class of carapace width of 11-12 cm and weight 70-90 g. The carapace width distribution shows that the carapace width capture >10 cm is 94%) **(Rahman *et al*., 2019**). Crab populations in coastal waters of the northern Persian Gulf, Iran dominated by adult crabs (carapace width >100 mm) from December to May, juvenile crabs are dominant from June to October (**Safaie *et al*., 2015**). The width of the crab carapace ranges from 60-150 mm (males) and 50-145 mm (females). The total weight males ranges to 48.0-275.50 g and females from 39.50 to 255.20 g **(Hosseini *et al*., 2012)**.

The length-weight relationship is one of the most important biological characters of fishes where weight of the fish increases as a function of its length (**Abeer *et al*, 2022**). **Fig (3)** exponent value (b) different to 3, this means the increase in carapace width is not as fast as the increase in weight. Coefficient b of 2.7344 shows growth allometrik negatif (P<0,05).

The same research was obtained **Muhsoni *et al* (2009)** in Bangkalan-Madura Waters, both males and females b<3. The same results were also obtained in the study **(Kanedi, 2020**) in the East Lampung Waters. The results are different from (**Ningrum, 2015**) which shows a value of b>3 (Allometrik positif) in the waters of Betahwalang and also at TPI Bulu, Jepara with growth pattern allometric positif **(Iksanti, 2022**).

Carapace width-total weight relationship showed positive allometric growth (b-value= 3.10)(**El-Kasheif, 2021**). **Afzaal (2018)** reports the carapace length-weight of separate sexes (male

Commented [AMT'S11]: Consistency in unit use..

Commented [AMT'S12R11]..... if the author use cm, convert mm to cm, to facilitate comparative discussion.

Commented [AMT'S13R11].....and vice versa if you use mm, convert cm to mm.

Commented [AMT'S14]: positive allometric

and female) of *P. pelagicus* were also calculated as $W = 0.967L^{2.91}$ (R2 = 0.874) and carapace width-weight were as $W = 0.8138L^{2.95}$ (R2 = 0.882).

Blue swimming crab male and female crabs found in Bone waters consists of 1-3 cohorts (**Kembaren *et al*., 2012**). The existence of cohort differences is suspected of subpopulation changes as a result of fishing pressure where fishing pressures of the sub population is decreasing (**Ernaningsih *et al*., 2022**).

Value of the growth parameter Von Bertalanffy predicting the relationship between Fish stock estimation and fisheries resource management **(Abeer *et al,* 2022).** The value of K indicates the growth rate blue swimming crab slow and takes a long time to reach its maximum length. Different growth coefficients in crustaceans can be due to several factors that is age, genetics, and seasons that affect water temperature (**Sara *et al*., 2016**).

Results of research on growth parameters of blue swimming crab (*P. pelagicus*) At different locations and times show different values (**Table. 2**).

Commented [AMT'S15]: at

Table 2. Growth parameter of blue swimming crabs (*P. pelagicus*) at different locations and times.

Total mortality rate (Z), natural mortality (M) dan fishing mortality (F) this study is lower than obtained by **Iksanti, *et al* (2022).** found a blue swimming crab landed at TPI Jepara at total mortality rate (Z) 2.99/year, natural mortality (M) 1,31/year dan fishing mortality (F) sebesar 1,68/year. Total mortality rate (Z), natural mortality (M) and fishing mortality (F) *of P. pelagicus* caught in Bardawil Wetland, Egypt 4.68/year, 2.25/year and 2.43/year respectively (**El-Betar, 2022**). The total mortality, natural mortality and capture mortality rates *of P. pelagicus* found in Egypt's Red Sea waters were 2,929/year, 1,285/year and 1,644/year, respectively (**El-Kasheif, 2021).** Fish species are more susceptible to activities fishing rather than deaths caused naturally by predation, competition, and disease (**Ofori-Danson *et al,* 2021**). The exploitation pressure of a fishing area can also be related to the parameters the dynamics of the fish population (**Ernaningsih *et al,* 2022**)

Commented [AMT'S16]: ,

The ratio of exploitation is very important to estimate the state of the stock is optimal, under exploitation or over exploitation (**Mehanna *et al.,* 2019**). The ratio of exploitation of *P. pelagicus* found in this study in Bawasalo waters Pangkep Regency of 0.55 indicates that exploitation ratio exceeds the optimal limit (MSY). The results of interviews with local fishermen that the capture of blue swimming crab the intensity is quite high where fishermen catch every day using several types of fishing gear. In the Tiworo Strait shows that the population of blue swimming crab in the category of moderate exploitation (**Permatahati *et al,* 2020**).

CONCLUSION

Distribution carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a

proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), the growth rate is low, the mortality rate of arrest is greater than the natural mortality rate and the exploitation ratio at limit of maksimal eksploitation (MSY) .

Commented [AMT'S17]: Maximum Sustainable Yield

REFERENCES

Abrenica, B. T. (2021). Stock Assessment of the Blue Swimming Crab *Portunus pelagicus* (Linnaeus, 1758) in Stock Enhancement Sites of Danajon Bank, Central Philippines. *Philippine Journal of Fisheries*, *28*(2), 210–227. https://doi.org/10.31398/tpjf/28.2.2020C0014

Afzaal, Z.; Kalhoro, M.; Buzdar, M.; Tariq, S., Shafi, M.; Nadeem, A.; Imran, S.; Saeed, F.; Sohail, M.; Hassan, R.; Haroon, A.; Shah, H. and Ahmed, I. (2018). Carapace length-weight and carapace width-weight relationship of *Portunus pelagicus* (Linnaeus, 1758) in Pakistani waters northern Arabian Sea. *Indian Journal of Geo-Marine Sciences*, *47*(4), 890–896. https://api.elsevier.com/content/abstract/scopus_id/85045961734

Agus, S. B.; Zulbainarni, N.; Sunuddin, A.; Subarno, T.; Nugraha, A. H.; Rahimah, I.; Alamsyah, A.; Rachmi, R. and Jihad. 2016. Spatial Distribution of blue swimming crabs (*Portunus pelagicus*) in the Eastern Season in the waters of Lancang Island, Thousand Islands. Indonesian Journal of Agricultural Sciences 21(3): 209-218.

Asiedu, B.; Amponsah, S. K. K.; Ofori-Danson, P. K. and Nunoo, F. K. E. (2023). Assessment of Exploited Population Dynamics of African Butter Catfish (Schilbe mystus) in Stratum VII of Lake Volta, Ghana. *Egyptian Journal of Aquatic Biology and Fisheries*, *27*(6), 153–168. https://doi.org/10.21608/ejabf.2023.328024

Beverton, R.J.H. and Holt, S.J. (1956). A review of methods for estimation of mortality rates in exploited fish populations with special reference to sources of bias in catch sampling. Rapp-v Reun. Cons Intern. Mer., 140:67-83.

El-Betar, T. A. (2022). Population structure and growth aspects of blue swimming crab, Portunus pelagicus, in Bardawil Wetland, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *26*(4), 885–903. https://doi.org/10.21608/ejabf.2022.255480

Ernaningsih.; Muh Jamal.; Hasnidar. And Hadijah. (2022). Population parameters and exploitation rates of tiger grouper (*Epinephelus fuscoguttatus*) in the Spermonde Islands of South Sulawesi Jurnal Bawal Widyariset Perikanan Tangkap. 14 (1) April 2022: 1-9. e-ISSN: 2502-6410.

El-Kasheif, M. A. (2021). Fishery Biology and Population Structure of the Blue Swimmer Crab, Portunus pelagicus, from the Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *25*(6), 269–283. https://doi.org/10.21608/EJABF.2021.211506

Hamid, A.; Wurdiatno, Y.; Batu, D. T. F. L. and Riani, E. (2015). Fecundity and Maturity of Rajungan Gonads (Portunus pelagicus) Females Incubate Eggs in Lasongko Bay,. Southeast Sulawesi. Bawal. 7(1), 43-50.

Hosseini, M.; Vazirizade, A.; Parsa, Y. and Mansori, A. (2012). Sex ratio, size distribution and seasonal abundance of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) in Persian Gulf Coasts, Iran. *World Applied Sciences Journal*, *17*(7), 919–925. https://www.scopus.com/inward/record.uri?eid=2-s2.0-84861959364&partnerID=40&md5=be3ff1b6f61442673ff55db3ab710992

Ihsan.; Eko, S. W.; Sugeng, H. W. and John, H. (2014). A Study of Biological Potential and Sustainability of Swimming Crab Population in the Waters of Pangkep Regency South Sulawesi Province. International Journal of Sciences: Basic and Applied Research. 16(1): 351-363.

Ihsan. (2018). Distribusi Ukuran Dan Pola Musim Penangkapan Rajungan (*Portunus pelagicus*) Di Perairan Kabupaten Pangkep, jurnal marine Fisheries, Hal: 73-83

Iksanti, R. M.; Redjeki, S. and Taufiq-Spj, N. (2022). Biological Aspects of blue swimming crab (*Portunus pelagicus*) Linnaeus, 1758 (Malacostraca: *Portunidae*). Reviewed from Morphometry and Gonad Maturity Level at TPI Bulu, Jepara. Journal of Marine Research, 11(3), 495-505.

I. Maiyza, S.; F. Mehanna, S. and A. El-karyoney, I. (2020). An evaluation for the exploitation level of Egyptian Marine Fisheries. *Egyptian Journal of Aquatic Biology and Fisheries*, *24*(7), 441–452. https://doi.org/10.21608/ejabf.2020.121292

Kanedi, M. M.; Rahardjo, P. and Maulita, M. (2020). Biological Aspects of blue swimming crab (*Portunus pelagicus*) in Coastal East Lampung Regency,Provinsi Lampung. Buletin Jalanidhitah Sarva Jivitam, 2(1), 49-56.

Kembaren, D.D.; Ernawati, T. and Suprapto. (2012). Biology and population parameters of crabs (*Portunus pelagicus*) in Bone Waters and surrounding areas. Jurnal Balai Penelitian Laut Jakarta. Vol 18 No. 4. Desember 2012 : 273-281 (In Indonesia)

Kusuma, B. T. dan Dewi. D. M. (2017). The Role of Fisheries Policies and Institutions in Management of Knitting (P*ortunus pelagicus*) So that it becomes a sustainable fishery Prosiding Simposium Nasional Krustasea 2017.

Lin, Y. J.; Grandcourt, E. M.; Rabaoui, L.; Maneja, R. H.; Qurban, M. A.; Al-Abdulkader, K. and Roa-Ureta, R. H. (2019). Comparative analysis of life history traits and trends of abundance in coral reefs of the orange-spotted grouper (Epinephelus coioides) from two regions of the Arabian Gulf. *ICES Journal of Marine Science*, *76*(4), 987–998. https://doi.org/10.1093/icesjms/fsz006

Mehanna, S. F.; Makkey, A. F. and Desouky, M. G. (2018). Growth, mortality and relative yield per recruit of the sharptooth catfish clarias gariepinus (Clariidae) in Lake Manzalah, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *22*(5), 65–72.

Mehanna, S. F.; Osman, Y. A. A.; Khalil, M. T. and Hassan, A. (2019). Age and growth, mortality and exploitation ratio of epinephelus summana (Forsskål, 1775) and cephalopholis argus (schneider, 1801) from the egyptian red sea coast, hurghada fishing area. *Egyptian Journal of Aquatic Biology and Fisheries*, *23*(4), 65–75.

Nurdin, M.; S, Ali.; S. A, dan Yanuarita. D. (2015). Mortalitas dan Laju Eksploitasi Rajungan (*Portunus pelagicus*) di Perairan Pulau Salemo Kabupaten Pangkajene Kepulauan. Jurnal Ipteks PSP 2:316-321.

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016a). Sex Ratio and Size at First Maturity Of Blue Swimming Crab (Portunus pelagicus) at Salemo Island, South Sulawesi.Ilmu Kelautan 21:17-22

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016b).. Size Distribution and Growth Pattern of Rajungan (*Portunus pelagicus*) on Salemo Island, Pangkajene Islands Regency. at the National Symposium on Marine Affairs and Fisheries III Makassar 7 Mei 2016. Page : 315-322

Ningrum, V. P.; Ghofar, A. and Ain, C. (2015). Biological Aspects Of Blue Swimmer Crab (*Portunus Pelagicus*) In Betahwalang Waters And Around. Saintek Perikanan: Indonesian Journal Of Fisheries Science And Technology, 11(1), 62-71.

Ofori-Danson, P.K.; Asiedu, B.; Alhassan, E.H.; Atsu, D.K. and Amponsah, S.K.K. (2021). Pre-Impoundment Fish Stock Assessment of the Black Volta: A Contribution to Fisheries Management of Bui Reservoir in Ghana. West African Journal of Applied Ecology, vol. 29(1): 35 – 48.

Pauly, D. (1979). Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries (No. 1; ICLARM, ICLARM Studies and Reviews). https://digitalarchive.worldfishcenter.org/ handle/ 20.500.12348/3693.

Pauly, D. (1980). On the interrelationship between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Intern.Explor. Mer., 39(2): 175-192.

Pauly, D. (1983). Some simple methods for the assessment of tropical fish stocks. FAO, Fish. Tech. Pap., 234: pp. http://www.fao.org/DOCREP/003/X6845E/ X684 5E00.HTM

Rahman, M. A.; Iranawati, F.; Sambah, A. B. and Wiadnya, D. G. R. (2019). Biological condition and carapace width frequency distribution of blue swimming crabs (Portunus pelagicus) in Gresik and Lamongan, East Java. *IOP Conference Series: Earth and Environmental Science*, *370*(1). https://doi.org/10.1088/1755-1315/370/1/012010

Permatahati, Y. I.; Bugis, N. N.; Sara, L. and Hasuba, T. F. (2020). Stock status of blue swimming crab (Portunus pelagicus Linnaeus, 1758) in Tiworo Strait waters, Southeast Sulawesi, Indonesia. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, *25*(2), 85–90. https://doi.org/10.14710/ik.ijms.25.2.85-90

Rabaoui, L. (2021). Distribution, abundance, and life history traits of the blue swimming crab Portunus segnis (Forskål, 1775) in the Saudi waters of the Arabian Gulf. *Regional Studies in Marine Science*, *46*. https://doi.org/10.1016/j.rsma.2021.101895

Russ, G.R. 1991. Coral Reef Fisheries: Effect and Yields. In:The Ecology of Fishes On Coral Reef. Ed. P.F Sale, pp.601-35. Academic Press Limited, Lonndon.

Safaie, M.; Shokri, M. R.; Kiabi, B. H. and Pazooki, J. (2015). Biomass, CPUE and size frequency distribution of blue swimming crab Portunus segnis (Forskal, 1775) in coastal waters of the northern Persian Gulf, Iran. *Journal of the Marine Biological Association of the United Kingdom*, *95*(4), 763–771. https://doi.org/10.1017/S0025315414001635

Sara, L. and Astuti, O. (2019). Status of blue swimming crab (Portunus pelagicus) population in southeast sulawesi waters, Indonesia. *AACL Bioflux*, *12*(5), 1909–1917. https://www.scopus.com/inward/record.uri?eid=2-s2.0-85075242107&partnerID=40&md5=b0dc93ae0c32e8fa7812b1e9b52d2e3a

Sparre, P. and Venema, S. C. (2005). Introduction to TropicalFish Stock Assesment (Part 1, p. 376)

Sujatha, K.; Shrikanya, K. V. L. and Iswarya Deepti, V.A. (2015). Species diversity and some aspects ofreproductive biology and life history of groupers(Pisces: Serranidae: Epinephelinae) off the centraleastern coast of India. Marine Biology Research,11(1), 18–33. https://doi.org/10.1080/17451000.2014.949271

Tirtadanu. (2019). Fishery, population parameters and exploitation status of blue swimming crab (Portunus pelagicus) in kwandang waters, indonesia. *AACL Bioflux*, *12*(4), 1323–1334. https://api.elsevier.com/content/abstract/scopus_id/85073378715

Then, A. Y.; Hoenig, J. M. and Huynh, Q. C. (2018). Estimating fishing and natural mortality rates, andcatchability coefficient, from a series of observationson mean length and fishing effort. ICES Journal ofMarine Science, 75(2), 610–620. https://doi.org/10.1093/icesjms/fsx177.

Wiyono, E. S. (2015). The dynamic of landing blue swimming crab (Portunus pelagicus) catches in Pangkajene Kepulauan, South Sulawesi, Indonesia. *AACL Bioflux*, *8*(2), 134–141. https://api.elsevier.com/content/abstract/scopus_id/84925943854



Fig (1). Map of research location *P. pelagicus* in Bawasalo waters

Commented [AMT'S1]: Provide "Map Legend" and source of map.




Fig (2). Carapace width distribution of blue swimming crabs (*P. pelagicus*) in Bawasalo Waters




Fig (3). Width-weight Relationship of blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig (4). Cohort groups by Bhattacharya's methode in Bawasalo Waters




Fig. (5). Growth curve of Von Bertalanffy blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig. (6). Growth curve of Von Bertalanffy blue swimming crabs (*P. pelagicus*) in Bawasalo Waters




Fig. (7). Probability of capture




Fig. (8). Yield per recruit, biomass per recruit of blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig. (9). Relative yield per recruit and relative biomass per recruit in relation to different exploitation ratio of *P.pelagicus* in Bawasalo Waters

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Manuscript ID EJABF-2403-4089

1 message

Ernaningsih Aras <ernaningsih.aras@umi.ac.id> Mon, Apr 15, 2024 at 7:36 PM
To: mtkhalil52@hotmail.com

dear Chief-editor
I am sending a journal manuscript which I have revised according to the Reviewer's suggestions with the title Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
regards.
Ernaningsih

Manuscript ID EJABF-2403-4089.doc
1203K

Egyptian Journal of Aquatic Biology & Fisheries
Zoology Department, Faculty of Science,
Ain Shams University, Cairo, Egypt.
ISSN 1110 – 6131
Vol. 28(2): … – … (2024)
www.ejabf.journals.ekb.eg



Population parameters and exploitation ratio of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi

Ernaningsih*[1], Danial[1], Syahrul[1], Moh.Tauhid Umar[2], Nursyahran[3]

[1-]**Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90231**
[2-]**Faculty of Marine Science and Fisheries, Hasanuddin University Makassar**
[3-]**Institute of Maritime Technology and Business Balikdiwa, Makassar 90245**
Corresponding Author : ernaningsih.aras@umi.ac.id

ARTICLE INFO
Article History:
Received:
Accepted:
Online:

Keywords:
Portunus pelagicus, growth, mortalities, exploitation ratio, Bawasalo Waters

ABSTRACT

A total of 1315 individuals were collected from Bawasalo Waters during the period March – August 2023 From the catch of fishermen with bubu naga fishing gear. This study aims to analyze population parameters and exploitation rate of Blue swimming crab (*Portunus pelagicus*) who was caught with a bubu naga in the waters of Bawasalo, Pangkep District, South Sulawesi.. The observed parameters include the size distribution of the carapace width , carapace width-weights relationship, Age group, Growth rate , total mortality rate, natural rate, fishing rate and exploitation ratio analyzed with the ELEFAN II program. Size carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), The Von Bertalanffy growth parameters were estimated to be K = 0,41 yr^{-1}, L maks = 15.10 cm, t0 = -0,723 yr, value total mortality (Z), natural mortality (M) and fishing mortality (F) were calculated as 2,70, 1.22, 1.48 yr^{-1} respectively. Exploitation rate (E = 0.55) indicates that the stock of *P. pelagicus* was slightly higher than that optimally exploited stock

INTRODUCTION

Blue swimming crab is widely distributed throughout Indonesia (**Kusuma dan Dewi, 2017**). In several regions in Indonesia, blue swimming crab is used as one of the people's fishery businesses including the Thousand Islands (**Agus *et al.*, 2016**),

Indexed in Scopus ELSEVIER DOAJ IUCAT

Lasongko Bay (**Hamid *et al*., 2015**;) and Pangkep Regency **(Ihsan *et al*., 2014**; **Nurdin *et al*., 2016**).

Pangkep Regency located in South Sulawesi has waters with considerable blue swimming crab (*P. pelagicus*) potential and became one of the important commodities. In Pangkep Regency itself, one of the blue swimming crab producing centers is on Salemo Island (**Nurdin *et al*., 2016b**). Total exports of blue swimming crab in 2020 reached 27,616,332 tons, with export value reaching USD 367,519,713 (**KKP, 2021**) is fourth in the total export value of Indonesian fishery products after shrimp, tuna and squid / cuttlefish / octopus.

The high export value indicates high production activities and exploitation of blue swimming crab, supported by open access fishing activities. Blue swimming crab (*P. pelagicus*) is currently under considerable pressure as the volume and size of the catch decreases (**Wiyono, 2015**). High fishing pressure from blue swimming crab (*P. pelagicus*) in some areas in Indonesia has led to reduced stocks (**Tirtadanu, 2019**). The catch and abundance of blue swimming crab on Salemo Island, Pangkep Regency has decreased and affected the income of fishermen (**Nurdin *et al*, 2015**). Fisheries resources have seriously declined due to overfishing, illegal fishing activities and pollution **(Lin *et al*., 2019) ; Mehanna *et al*., 2018**) **; Asiedu *et al*., 2023**).

Biological information and population dynamics are very important for stock estimation (**Then *et al*, 2018**). Information about biological aspects is a major factor for management purposes (**Sujatha *et al*, 2015**). For management purposes, information on fisheries biology and population structure is required (**El-Kasheif, 2021**). The sustainability of the blue swimming crab requires population biology information and estimating fishery stocks as a scientific basis study for the availability of sustainable resources **(I. Maiyza *et al*., 2020; Froese *et al*, 2011**). Research related to blue swimming crab in Pangkep Regency has been made regarding the arrest (**Ihsan, 2018).** Bawasalo waters, is one of the blue swimming crab fishing areas in Pangkep Regency, However a study of population parameters and exploitation ratio of blue swimming crab (*P. pelagicus*) those caught with bubu naga in Bawasalo waters have not been found. Based on this, it is important to conduct research related to population parameters and exploitation rates which include the distribution of carapace width, carapace width-weight relationship, age group, growth rate, mortality rate dan rate of exploitation as one of the important studies in the sustainability of blue swimming crab resources.

MATERIALS AND METHODS

Total of 1159 blue swimming crabs (*P. pelagicus*) randomly drawn from March 2023 to August 2023 is the catch of bubu naga originating from Bawasalo Waters, Segeri District, Pangkep Regency (**Fig. 1**). Each sample of blue swimming crab measured carapace width (CW) in cm (The width of the blue swimming crab is measured from the longest lateral spines on the sides of its body) and weight (W) in g.




Figure 1. Map of research location *P. pelagicus* in Bawasalo waters

Length frequency data was analyzed by Bhattacharya's method (**Bhattacharya, 1967**) that was involved in the FiSAT II software package (**Gayanilo, 2005**).

The total length and body weight of fish were used for the length-weight relationship. The length-weight relationship was analyzed using the equation below (**Ricker, 1975**):

$$W = aL^b$$

Where, W = Total body weight of the fish (g); L = Total carapace width (cm); "a" = Regression constant, and "b" = Koefisien regresi; Value "b" provide information about the type of fish growth. The degree of association between length-weight variables is calculated by the correlation coefficient (R2), and the degree of its statistical significance is assessed following the method indicated by **Santos *et al.* (2002)**, as follows:

$$\text{Log } W = \text{Log } a + b \text{ Log } TL$$

If b = 3, The growth is isometric, b ≠ 3, The growth is allometric (allometric negative if b < 3 and allometric positive if b > 3).

Growth parameters follow the growth function of Von Bertalanffy (VBGF) including growth rate (K) and asymptotic length (L∞) with equations.

$$L_t = L_\infty\left(1 - e^{-K(t - t_0)}\right)$$

Where, Lt is the average length at time (or age) t, L maks; is the asymptotic mean length; K is the coefficient of growth rate (per year), t; represents time or age. The theoretical age at zero length (t0) follows the equation shown by **Pauly (1979)**, as follows:

Log 10 (-t0) = -0.3922 – 0.2752 log 10 L∞ - 1.038log 10 K

Estimation of total mortality (Z) was carried out by the length converted catch curve method **(Pauly, 1983)**. Natural mortality rate **(M) = Log M = -0.0066 – 0.279 log L∞ + 0.6543 log K + 0.4634 log T (Pauly, 1980),** with a flat temperature of 30^0 C. Fishing mortality rate (F)= Z-M. Exploitation rate using equations **Gulland (1971)**, **E = F/Z**

Long was first caught allegedly based on probabilitas fishing cumulative against the middle of the width class. From the resulting curve, Long first caught (Lc50) which is defined as the average total length where 50% of the fish are caught, taken according to the cumulative probability at 50%. This procedure follows the method by **Beverton dan Holt (1956**) dan **Sparre dan Venema (1998)**, as follows :

Lc50 = [TL – K(TL∞ – TL‟)]/ Z

Where , Lc50 is the first length, and T times caught where L 'is the average length of fish in the catch sample. Moreover, length at 75 and 95% fishing correlates with cumulative probability at 75 and 95%, respectively, as shown by **Pauly (1983).**

The result per recruit (Y / R) is determined according to the formula developed by **Beverton dan Holt (1957**), dan hasil relatif per rekrut (Y / R0) dihitung menurut **Pauly dan Soriano (1986).**

Data analysis

The FiSAT II tool was used to assess the population parameters, such asymptotic length (L∞), growth rate (K), natural mortality rate (M), fishing mortality rate (F), exploitation ratio (E), and relative yield per recruit indicators of samples from the assessed fish species encountered during the study period (**Gayanilo *et al*. 2005)**

RESULTS

Carapace width distribution

Carapace width distribution of blue swimming crab carapace (*P. pelagicus*) during the study from March 2023-August 2023, Who was caught with the bubu naga in Bawasalo waters Pangkep Regency to 1315 individuals on the group width carapace 6,70 cm – 13,80 cm, with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) (**Fig. 2**).




Fig (2). Carapace width distribution of blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

Carapace Width-weight relationship

The carapace width and weight relationship for males, females and combined sexes of *P. pelagicus* was described by equation as : $W= 0.1052L^{2.7245}$ (R2 = 7478), **(Fig 3**).




Fig (3). Width-weight Relationship of blue swimming crab (*P. pelagicus*) in Bawasalo waters

Age Group

Cohort analysis with Battacahrya's method *P. Pelagicus Consists of 3 age groups* is shown in **Fig (4).** Three cohort group were identified at length 8.66 sd 0.67, 10.27 sd 0.93, 12.16 sd 0.21 cm




Fig (4). Cohort groups by Bhattacharya's methode in Bawasalo Waters

Growth parameters

Estimated values of growth parameters were analyzed using the FISAT II program, The results of the analysis of the growth parameters estimation with the formula von Bertalannffy of blue swimming crab *P. pelagicus* including maximum length (L∞), growth coefficient (K) and age at birth (t0) that is L∞ = 15.1 cm, K = 0.41 per month and t0 = -0.48 **(Fig 5).**




Fig. (5). Growth curve of Von Bertalanffy blue swimming crab (*P. pelagicus*) in Bawasalo waters

Moralities and exploitation ratio

Mortality analysis based length-converted catch curve (**Fig. 6**) obtained total mortality rate, natural mortality, fishing mortality dan exploitation ratio blue swimming crab (*P. pelagicus*).




Fig. (6). Growth curve of Von Bertalanffy blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

The total mortality (Z) was 2.70 yr^{-1}, natural mortality (M) and fishing mortality (F) was 1.22 yr^{-1} and 1.48 yr^{-1} and the exploitation ratio was 0.55 (**Table 1**).

Table 1. Value of population paramters blue swimming crab (*P. pelagicus*) in Bawasalo waters

Parameters	Value
Total mortality (Z)	2.70 yr-1
Natural mortality (M)	1.22 yr-1
Fishing mortality (F)	1.48 yr-1
Length at first capture (Lc)	8.56 cm
Length of recruitment (Lr)	8.22
Age at first capture (tc)	1.56
Age of recruitment (tr)	1.43
Exploitation ratio (E cur.)	0.55
The yield per recruit (Y/R)	20.6g
Biomass per recruit (B/R)	12.3g
The relative yield per recruit (Y/R')	0.009
Lc/L∞	0.567
M/K	2.976

Length at first capture

The length first capture (Lc50) is 10.92 cm. Length of catch probability 25% (Lc25) and 75% (Lc75) as 10.53 cm dan 11.22 cm **(Fig. 7).**




Fig. (7). Probability of capture

Yield per recruit (Y/R), and relative yield per recruit (Y'/R)

The yield per recruit (Y/R) for P. pelagicus was 20.6 gr pada current level mortality (F cur) 4.8 yr^{-1} (**Fig 8**). Relative yield per Y'/R recruit for different exploitation ratio (**Fig 9)**. From this curve, the maximum Y'/R is found at the exploitation ratio (Emax) = 0,50. The exploitation ratio E = 0.3.. would give the maximum Y/R' results in the protection of about 50% of the stock biomass




Fig. (8). Yield per recruit, biomass per recruit of blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig. (9). Relative yield per recruit and relative biomass per recruit in relation to different exploitation ratio of *P.pelagicus* in Bawasalo Waters

DISCUSSION

Carapace width distribution of blue swimming crab (*P. pelagicus*) during the study that is 6,70 cm – 13,80 cm, where the size of the carapace width < 10 cm as 66,69% and carapace width >10 cm as 33,31%**.** Research results **Ihsan (2018)** in Pangkep's waters finding the dominant class interval males blue swimming crab on size 112-123 mm with a frequency of 265 heads (31,77%) and blue swimming crab females at intervals of classes 100-111 mm with a frequency of 223 heads (26,74%). Research results **Abdul Hamid (2019**) in Lasongko Bay shows carapace width on male blue swimming crab ranges from 4.1 cm -12.4 cm. Size of *P. pelagicus* obtained in Kwandang waters ranges from 7.0 – 17.5 cm (**Tirtadanu *et al.*, 2019**). Blue swimming crab catches obtained in Gresik and Lamongan dominated by the size class of carapace width of 11-12 cm and weight 70-90 g. The carapace width distribution shows that the carapace width capture >10 cm is 94%) **(Rahman *et al.*, 2019**). Crab populations in coastal waters of the northern Persian Gulf, Iran dominated by adult crabs (carapace width >100 mm) from December to May, juvenile crabs are dominant from June to October (**Safaie *et al.*, 2015**). The width of the crab carapace ranges from 60-150 mm (males) and 50-145 mm (females). The total weight males ranges to 48.0-275.50 g and females from 39.50 to 255.20 g **(Hosseini *et al.*, 2012)**. The length-weight relationship is one of the most important biological characters of fishes where weight of the fish increases as a function of its length (**Abeer *et al*, 2022**). **Fig (3)** exponent value (b) different to 3, this means the increase in carapace width is not as fast as the increase in weight. Coefficient b of 2.7344 shows growth allometrik negatif (P<0,05).

The same research was obtained **Muhsoni *et al* (2009)** in Bangkalan-Madura Waters, both males and females b<3. The same results were also obtained in the study **(Kanedi, 2020**) in the East Lampung Waters. The results are different from (**Ningrum, 2015**) which shows a value of b>3 (Allometrik positif) in the waters of Betahwalang and also at TPI Bulu, Jepara with growth pattern allometric positif **(Iksanti, 2022**). Carapace width-total weight relationship showed positive allometric growth (b-value= 3.10)(**El-Kasheif, 2021**). **Afzaal (2018)** reports the carapace length-weight of separate sexes (male and female) of *P. pelagicus* were also calculated as $W = 0.967L^{2.91}$ (R2 = 0.874) and carapace width-weight were as $W = 0.8138L^{2.95}$ (R2 = 0.882).

Blue swimming crab male and female crabs found in Bone waters consists of 1-3 cohorts (**Kembaren *et al.*, 2012**). The existence of cohort differences is suspected of subpopulation changes as a result of fishing pressure where fishing pressures of the sub population is decreasing (**Ernaningsih *et al.*, 2022**).

Value of the growth parameter Von Bertalanffy predicting the relationship between Fish stock estimation and fisheries resource management **(Abeer *et a*l, 2022).** The value of K indicates the growth rate blue swimming crab slow and takes a long time to reach its maximum length. Different growth coefficients in crustaceans can be due to several factors that is age, genetics, and seasons that affect water temperature (**Sara *et al.*,**

2016). Results of research on growth parameters of blue swimming crab (*P. pelagicus*) At different locations and times show different values (**Table. 2**).

Table 2. Growth parameter of blue swimming crabs (*P. pelagicus*) at different locations and times.

Researcher/year	Location	K (yr $^{-1}$)	Lmaks (cm)
Kembaren/2012	Bone Bay	1,27	15.9
Sara/2017	Southeast Sulawesi Waters	0.55	16.9
Tirtadanu/2019	Indonesia	1,24	-
Abrenica/2021	Philipines	1,30	21,65
Rabaoui/2021	Saudi Arabian	2,12	-
Kasheif, 2021	Red Sea, Egypt	0,41	21.19
El-Betar/2022	Egypt	1,40	16.60
The present study/2023	Bawasalo, Pangkep	0,41	15.10

Total mortality rate (Z), natural mortality (M) dan fishing mortality (F) this study is lower than obtained by **Iksanti, *et al* (2022).** found a blue swimming crab landed at TPI Jepara at total mortality rate (Z) 2.99/year, natural mortality (M) 1,31/year dan fishing mortality (F) sebesar 1,68/year. Total mortality rate (Z), natural mortality (M) and fishing mortality (F) *of P. pelagicus* caught in Bardawil Wetland, Egypt 4.68/year, 2.25/year and 2.43/year respectively (**El-Betar, 2022**). The total mortality, natural mortality and capture mortality rates *of P. pelagicus* found in Egypt's Red Sea waters were 2,929/year, 1,285/year and 1,644/year, respectively (**El-Kasheif, 2021).** Fish species are more susceptible to activities fishing rather than deaths caused naturally by predation, competition, and disease (**Ofori-Danson *et al,* 2021**). The exploitation pressure of a fishing area can also be related to the parameters the dynamics of the fish population (**Ernaningsih *et al,* 2022**).

The ratio of exploitation is very important to estimate the state of the stock is optimal, under exploitation or over exploitation (**Mehanna *et al.,* 2019**). The ratio of exploitation of *P. pelagicus* found in this study in Bawasalo waters Pangkep Regency of 0.55 indicates that exploitation ratio exceeds the optimal limit (MSY). The results of interviews with local fishermen that the capture of blue swimming crab the intensity is quite high where fishermen catch every day using several types of fishing gear. In the Tiworo Strait shows that the population of blue swimming crab in the category of moderate exploitation (**Permatahati *et al,* 2020**).

CONCLUSION

Distribution carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), the growth rate is low, the mortality rate of arrest is greater than the natural mortality rate and the exploitation ratio at limit of maksimal eksploitation (MSY) .

REFERENCES

Abrenica, B. T. (2021). Stock Assessment of the Blue Swimming Crab *Portunus pelagicus* (Linnaeus, 1758) in Stock Enhancement Sites of Danajon Bank, Central Philippines. *Philippine Journal of Fisheries*, *28*(2), 210–227. https://doi.org/10.31398/tpjf/28.2.2020C0014

Afzaal, Z.; Kalhoro, M.; Buzdar, M.; Tariq, S., Shafi, M.; Nadeem, A.; Imran, S.; Saeed, F.; Sohail, M.; Hassan, R.; Haroon, A.; Shah, H. and Ahmed, I. (2018). Carapace length-weight and carapace width-weight relationship of *Portunus pelagicus* (Linnaeus, 1758) in Pakistani waters northern Arabian Sea. *Indian Journal of Geo-Marine Sciences*, *47*(4), 890–896. https://api.elsevier.com/content/abstract/scopus_id/85045961734

Agus, S. B.; Zulbainarni, N.; Sunuddin, A.; Subarno, T.; Nugraha, A. H.; Rahimah, I.; Alamsyah, A.; Rachmi, R. and Jihad. 2016. Spatial Distribution of blue swimming crabs (*Portunus pelagicus*) in the Eastern Season in the waters of Lancang Island, Thousand Islands. Indonesian Journal of Agricultural Sciences 21(3): 209-218.

Asiedu, B.; Amponsah, S. K. K.; Ofori-Danson, P. K. and Nunoo, F. K. E. (2023). Assessment of Exploited Population Dynamics of African Butter Catfish (Schilbe mystus) in Stratum VII of Lake Volta, Ghana. *Egyptian Journal of Aquatic Biology and Fisheries*, *27*(6), 153–168. https://doi.org/10.21608/ejabf.2023.328024

Beverton, R.J.H. and Holt, S.J. (1956). A review of methods for estimation of mortality rates in exploited fish populations with special reference to sources of bias in catch sampling. Rapp-v Reun. Cons Intern. Mer., 140:67-83.

El-Betar, T. A. (2022). Population structure and growth aspects of blue swimming crab, Portunus pelagicus, in Bardawil Wetland, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *26*(4), 885–903. https://doi.org/10.21608/ejabf.2022.255480

Ernaningsih.; Muh Jamal.; Hasnidar. And Hadijah. (2022). Population parameters and exploitation rates of tiger grouper (*Epinephelus fuscoguttatus*) in the Spermonde Islands of South Sulawesi Jurnal Bawal Widyariset Perikanan Tangkap. 14 (1) April 2022: 1-9. e-ISSN: 2502-6410.

El-Kasheif, M. A. (2021). Fishery Biology and Population Structure of the Blue Swimmer Crab, Portunus pelagicus, from the Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *25*(6), 269–283. https://doi.org/10.21608/EJABF.2021.211506

Hamid, A.; Wurdiatno, Y.; Batu, D. T. F. L. and Riani, E. (2015). Fecundity and Maturity of Rajungan Gonads (Portunus pelagicus) Females Incubate Eggs in Lasongko Bay,. Southeast Sulawesi. Bawal. 7(1), 43-50.

Hosseini, M.; Vazirizade, A.; Parsa, Y. and Mansori, A. (2012). Sex ratio, size distribution and seasonal abundance of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) in Persian Gulf Coasts, Iran. *World Applied Sciences Journal*, *17*(7), 919–925. https://www.scopus.com/inward/record.uri?eid=2-s2.0-84861959364&partnerID=40&md5=be3ff1b6f61442673ff55db3ab710992

Ihsan.; Eko, S. W.; Sugeng, H. W. and John, H. (2014). A Study of Biological Potential and Sustainability of Swimming Crab Population in the Waters of Pangkep Regency South Sulawesi Province. International Journal of Sciences: Basic and Applied Research. 16(1): 351-363.

Ihsan. (2018). Distribusi Ukuran Dan Pola Musim Penangkapan Rajungan (*Portunus pelagicus*) Di Perairan Kabupaten Pangkep, jurnal marine Fisheries, Hal: 73-83

Iksanti, R. M.; Redjeki, S. and Taufiq-Spj, N. (2022). Biological Aspects of blue swimming crab (*Portunus pelagicus*) Linnaeus, 1758 (Malacostraca: *Portunidae*). Reviewed from Morphometry and Gonad Maturity Level at TPI Bulu, Jepara. Journal of Marine Research, 11(3), 495-505.

I. Maiyza, S.; F. Mehanna, S. and A. El-karyoney, I. (2020). An evaluation for the exploitation level of Egyptian Marine Fisheries. *Egyptian Journal of Aquatic Biology and Fisheries*, *24*(7), 441–452. https://doi.org/10.21608/ejabf.2020.121292

Kanedi, M. M.; Rahardjo, P. and Maulita, M. (2020). Biological Aspects of blue swimming crab (*Portunus pelagicus*) in Coastal East Lampung Regency,Provinsi Lampung. Buletin Jalanidhitah Sarva Jivitam, 2(1), 49-56.

Kembaren, D.D.; Ernawati, T. and Suprapto. (2012). Biology and population parameters of crabs (*Portunus pelagicus*) in Bone Waters and surrounding areas. Jurnal Balai Penelitian Laut Jakarta. Vol 18 No. 4. Desember 2012 : 273-281 (In Indonesia)

Kusuma, B. T. dan Dewi. D. M. (2017). The Role of Fisheries Policies and Institutions in Management of Knitting (P*ortunus pelagicus*) So that it becomes a sustainable fishery Prosiding Simposium Nasional Krustasea 2017.

Lin, Y. J.; Grandcourt, E. M.; Rabaoui, L.; Maneja, R. H.; Qurban, M. A.; Al-Abdulkader, K. and Roa-Ureta, R. H. (2019). Comparative analysis of life history traits and trends of abundance in coral reefs of the orange-spotted grouper (Epinephelus coioides) from two regions of the Arabian Gulf. *ICES Journal of Marine Science*, *76*(4), 987–998. https://doi.org/10.1093/icesjms/fsz006

Mehanna, S. F.; Makkey, A. F. and Desouky, M. G. (2018). Growth, mortality and relative yield per recruit of the sharptooth catfish clarias gariepinus (Clariidae) in Lake Manzalah, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *22*(5), 65–72.

Mehanna, S. F.; Osman, Y. A. A.; Khalil, M. T. and Hassan, A. (2019). Age and growth, mortality and exploitation ratio of epinephelus summana (Forsskål, 1775) and cephalopholis argus (schneider, 1801) from the egyptian red sea coast, hurghada fishing area. *Egyptian Journal of Aquatic Biology and Fisheries*, *23*(4), 65–75.

Nurdin, M.; S, Ali.; S. A, dan Yanuarita. D. (2015). Mortalitas dan Laju Eksploitasi Rajungan (*Portunus pelagicus*) di Perairan Pulau Salemo Kabupaten Pangkajene Kepulauan. Jurnal Ipteks PSP 2:316-321.

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016a). Sex Ratio and Size at First Maturity Of Blue Swimming Crab (Portunus pelagicus) at Salemo Island, South Sulawesi.Ilmu Kelautan 21:17-22

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016b).. Size Distribution and Growth Pattern of Rajungan (*Portunus pelagicus*) on Salemo Island, Pangkajene Islands Regency. at the National Symposium on Marine Affairs and Fisheries III Makassar 7 Mei 2016. Page : 315-322

Ningrum, V. P.; Ghofar, A. and Ain, C. (2015). Biological Aspects Of Blue Swimmer Crab (*Portunus Pelagicus*) In Betahwalang Waters And Around. Saintek Perikanan: Indonesian Journal Of Fisheries Science And Technology, 11(1), 62-71.

Ofori-Danson, P.K.; Asiedu, B.; Alhassan, E.H.; Atsu, D.K. and Amponsah, S.K.K. (2021). Pre-Impoundment Fish Stock Assessment of the Black Volta: A Contribution to Fisheries Management of Bui Reservoir in Ghana. West African Journal of Applied Ecology, vol. 29(1): 35 – 48.

Pauly, D. (1979). Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries (No. 1; ICLARM, ICLARM Studies and Reviews). https://digitalarchive.worldfishcenter.org/ handle/ 20.500.12348/3693.

Pauly, D. (1980). On the interrelationship between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Intern.Explor. Mer., 39(2): 175-192.

Pauly, D. (1983). Some simple methods for the assessment of tropical fish stocks. FAO, Fish. Tech. Pap., 234: pp. http://www.fao.org/DOCREP/003/X6845E/ X684 5E00.HTM

Rahman, M. A.; Iranawati, F.; Sambah, A. B. and Wiadnya, D. G. R. (2019). Biological condition and carapace width frequency distribution of blue swimming crabs (Portunus pelagicus) in Gresik and Lamongan, East Java. *IOP Conference Series: Earth and Environmental Science*, *370*(1). https://doi.org/10.1088/1755-1315/370/1/012010

Permatahati, Y. I.; Bugis, N. N.; Sara, L. and Hasuba, T. F. (2020). Stock status of blue swimming crab (Portunus pelagicus Linnaeus, 1758) in Tiworo Strait waters, Southeast Sulawesi, Indonesia. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, *25*(2), 85–90. https://doi.org/10.14710/ik.ijms.25.2.85-90

Rabaoui, L. (2021). Distribution, abundance, and life history traits of the blue swimming crab Portunus segnis (Forskål, 1775) in the Saudi waters of the Arabian Gulf. *Regional Studies in Marine Science*, *46*. https://doi.org/10.1016/j.rsma.2021.101895

Russ, G.R. 1991. Coral Reef Fisheries: Effect and Yields. In:The Ecology of Fishes On Coral Reef. Ed. P.F Sale, pp.601-35. Academic Press Limited, Lonndon.

Safaie, M.; Shokri, M. R.; Kiabi, B. H. and Pazooki, J. (2015). Biomass, CPUE and size frequency distribution of blue swimming crab Portunus segnis (Forskal, 1775) in coastal waters of the northern Persian Gulf, Iran. *Journal of the Marine Biological Association of the United Kingdom*, *95*(4), 763–771. https://doi.org/10.1017/S0025315414001635

Sara, L. and Astuti, O. (2019). Status of blue swimming crab (Portunus pelagicus) population in southeast sulawesi waters, Indonesia. *AACL Bioflux*, *12*(5), 1909–1917. https://www.scopus.com/inward/record.uri?eid=2-s2.0-85075242107&partnerID=40&md5=b0dc93ae0c32e8fa7812b1e9b52d2e3a

Sparre, P. and Venema, S. C. (2005). Introduction to TropicalFish Stock Assesment (Part 1, p. 376)

Sujatha, K.; Shrikanya, K. V. L. and Iswarya Deepti, V.A. (2015). Species diversity and some aspects ofreproductive biology and life history of groupers(Pisces: Serranidae: Epinephelinae) off the centraleastern coast of India. Marine Biology Research,11(1), 18–33. https://doi.org/10.1080/17451000.2014.949271

Tirtadanu. (2019). Fishery, population parameters and exploitation status of blue swimming crab (Portunus pelagicus) in kwandang waters, indonesia. *AACL Bioflux*, *12*(4), 1323–1334. https://api.elsevier.com/content/abstract/scopus_id/85073378715

Then, A. Y.; Hoenig, J. M. and Huynh, Q. C. (2018). Estimating fishing and natural mortality rates, andcatchability coefficient, from a series of observationson mean length and fishing effort. ICES Journal ofMarine Science, 75(2), 610–620. https://doi.org/10.1093/icesjms/fsx177.

Wiyono, E. S. (2015). The dynamic of landing blue swimming crab (Portunus pelagicus) catches in Pangkajene Kepulauan, South Sulawesi, Indonesia. *AACL Bioflux*, *8*(2), 134–141. https://api.elsevier.com/content/abstract/scopus_id/84925943854

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

4089

3 messages

Magdy Khalil <mtkhalil52@hotmail.com> Tue, Apr 16, 2024 at 6:38 AM
To: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Dear Authors,
Please revise carefully the attached version of your article before Language editing and publishing, especially the references, to match that of the Journal;

Example
Van der Geer, J.; Hanraads, J.A.J. ; Khalil, M. and Lupton, R.A. (2010). The art of writing a scientific article. J. Sci. Commun., 163**:** 51–59.

Correct the chronological order of the references inside the text;

Please Send the cost of publication ($160) to Magdy Tawfik Khalil Ibrahim
Commercial International Bank (CIB)
Address: Thawra Branch- Cairo, Egypt
Account No. 100016289684
SWIFT CODE: CIBEEGCX081
IBAN: EG450010000600000100016289684
Beneficial address: Faculty of Science, Ain Shams University, Cairo, Egypt
Please write your full name in the receipt of the Bank

From: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>
Sent: Monday, April 15, 2024 12:36 PM
To: mtkhalil52@hotmail.com <mtkhalil52@hotmail.com>
Subject: Manuscript ID EJABF-2403-4089

dear Chief-editor
I am sending a journal manuscript which I have revised according to the Reviewer's suggestions with the title Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
regards.
Ernaningsih

4089.doc
1097K

Ernaningsih Aras <ernaningsih.aras@umi.ac.id> Wed, Apr 17, 2024 at 10:16 AM
To: Magdy Khalil <mtkhalil52@hotmail.com>

Dear chief editor
I sent the revised journal as directed (title: Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
and I send proof of journal payment (ID EJABF-2403-4089)
Yours faithfully
Ernaningsih
[Quoted text hidden]

2 attachments

Manuscript ID EJABF-2403-4089.doc
1205K

payment (Ernaningsih).pdf
216K

Ernaningsih Aras <ernaningsih.aras@umi.ac.id> Tue, Apr 23, 2024 at 7:15 PM
To: Magdy Khalil <mtkhalil52@hotmail.com>

Dear
please provide information on the progress of my journal (title: Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
Yours faithfully
Ernaningsih
[Quoted text hidden]

Egyptian Journal of Aquatic Biology & Fisheries
Zoology Department, Faculty of Science,
Ain Shams University, Cairo, Egypt.
ISSN 1110 – 6131
Vol. 28(2): … – … (2024)
www.ejabf.journals.ekb.eg



Population parameters and exploitation ratio of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi

Ernaningsih*[1], Danial[1], Syahrul[1], Moh.Tauhid Umar[2], Nursyahran[3]

[1]-**Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90245**
[2]-**Faculty of Marine Science and Fisheries, Hasanuddin University Makassar**
[3]-**Institute of Maritime Technology and Business Balikdiwa, Makassar 90245**
***Corresponding Author : ernaningsih.aras@umi.ac.id**

ARTICLE INFO
Article History:
Received:
Accepted:
Online:

Keywords:
Portunus pelagicus, growth, mortalities, exploitation ratio, Bawasalo Waters

ABSTRACT

A total of 1315 individuals were collected from Bawasalo Waters during the period March – August 2023 From the catch of fishermen with bubu naga fishing gear. This study aims to analyze population parameters and exploitation rate of Blue swimming crab (*Portunus pelagicus*) who was caught with a bubu naga in the waters of Bawasalo, Pangkep District, South Sulawesi.. The observed parameters include the size distribution of the carapace width , carapace width-weights relationship, Age group, Growth rate , total mortality rate, natural rate, fishing rate and exploitation ratio analyzed with the ELEFAN II program. Size carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), The Von Bertalanffy growth parameters were estimated to be K = 0,41 yr^{-1}, L maks = 15.10 cm, t0 = -0,723 yr, value total mortality (Z), natural mortality (M) and fishing mortality (F) were calculated as 2,70, 1.22, 1.48 yr^{-1} respectively. Exploitation rate (E = 0.55) indicates that the stock of *P. pelagicus* was slightly higher than that optimally exploited stock

INTRODUCTION

Blue swimming crab is widely distributed throughout Indonesia (**Kusuma dan Dewi, 2017**). In several regions in Indonesia, blue swimming crab is used as one of the people's fishery businesses including the Thousand Islands (**Agus *et al.*, 2016**), Lasongko Bay (**Hamid *et al.*, 2015**;) and Pangkep Regency **(Ihsan *et al.*, 2014**; **Nurdin *et al.*, 2016**).

Pangkep Regency located in South Sulawesi has waters with considerable blue swimming crab (*P. pelagicus*) potential and became one of the important commodities. In Pangkep Regency itself, one of the blue swimming crab producing centers is on

Indexed in Scopus ELSEVIER DOAJ IUCAT

Salemo Island (**Nurdin *et al.*, 2016b**). Total exports of blue swimming crab in 2020 reached 27,616,332 tons, with export value reaching USD 367,519,713 (**KKP, 2021**) is fourth in the total export value of Indonesian fishery products after shrimp, tuna and squid / cuttlefish / octopus.

The high export value indicates high production activities and exploitation of blue swimming crab, supported by open access fishing activities. Blue swimming crab (*P. pelagicus*) is currently under considerable pressure as the volume and size of the catch decreases (**Wiyono, 2015**). High fishing pressure from blue swimming crab (*P. pelagicus*) in some areas in Indonesia has led to reduced stocks (**Tirtadanu, 2019**). The catch and abundance of blue swimming crab on Salemo Island, Pangkep Regency has decreased and affected the income of fishermen (**Nurdin *et al*, 2015**). Fisheries resources have seriously declined due to overfishing, illegal fishing activities and pollution **(Lin *et al*., 2019) ; Mehanna *et al*., 2018**) **; Asiedu *et al*., 2023**).

Biological information and population dynamics are very important for stock estimation (**Then *et al*, 2018**). Information about biological aspects is a major factor for management purposes (**Sujatha *et al*, 2015**). For management purposes, information on fisheries biology and population structure is required (**El-Kasheif, 2021**). The sustainability of the blue swimming crab requires population biology information and estimating fishery stocks as a scientific basis study for the availability of sustainable resources **(I. Maiyza *et al*., 2020; Froese *et al*, 2011**). Research related to blue swimming crab in Pangkep Regency has been made regarding the arrest (**Ihsan, 2018).** Bawasalo waters, is one of the blue swimming crab fishing areas in Pangkep Regency, However a study of population parameters and exploitation ratio of blue swimming crab (*P. pelagicus*) those caught with bubu naga in Bawasalo waters have not been found. Based on this, it is important to conduct research related to population parameters and exploitation rates which include the distribution of carapace width, carapace width-weight relationship, age group, growth rate, mortality rate dan rate of exploitation as one of the important studies in the sustainability of blue swimming crab resources.

MATERIALS AND METHODS

Total of 1159 blue swimming crabs (*P. pelagicus*) randomly drawn from March 2023 to August 2023 is the catch of bubu naga originating from Bawasalo Waters, Segeri District, Pangkep Regency (**Fig. 1**). Each sample of blue swimming crab measured carapace width (CW) in cm (The width of the blue swimming crab is measured from the longest lateral spines on the sides of its body) and weight (W) in g.




Figure 1. Map of research location *P. pelagicus* in Bawasalo waters

Length frequency data was analyzed by Bhattacharya's method (**Bhattacharya, 1967**) that was involved in the FiSAT II software package (**Gayanilo, 2005**).

The total length and body weight of fish were used for the length-weight relationship. The length-weight relationship was analyzed using the equation below (**Ricker, 1975**):

$$W = aL^b$$

Where, W = Total body weight of the fish (g); L = Total carapace width (cm); "a" = Regression constant, and "b" = Koefisien regresi; Value "b" provide information about the type of fish growth. The degree of association between length-weight variables is calculated by the correlation coefficient (R2), and the degree of its statistical significance is assessed following the method indicated by **Santos *et al*. (2002)**, as follows:

$$\text{Log } W = \text{Log } a + b \text{ Log } TL$$

If b = 3, The growth is isometric, b ≠ 3, The growth is allometric (allometric negative if b < 3 and allometric positive if b > 3).

Growth parameters follow the growth function of Von Bertalanffy (VBGF) including growth rate (K) and asymptotic length (L∞) with equations.

$$L_t = L_\infty\left(1 - e^{-K(t-t_0)}\right)$$

Where, Lt is the average length at time (or age) t, L maks; is the asymptotic mean length; K is the coefficient of growth rate (per year), t; represents time or age. The theoretical age at zero length (t0) follows the equation shown by **Pauly (1979)**, as follows:

$$\text{Log } 10 (-t0) = -0.3922 - 0.2752 \log 10 L\infty - 1.038 \log 10 K$$

Estimation of total mortality (Z) was carried out by the length converted catch curve method **(Pauly, 1983)**. Natural mortality rate **(M) = Log M = -0.0066 – 0.279 log L∞ +**

0.6543 log K + 0.4634 log T (Pauly, 1980), with a flat temperature of 30^0 C. Fishing mortality rate (F)= Z-M. Exploitation rate using equations **Gulland (1971)**, **E = F/Z**
Long was first caught allegedly based on probabilitas fishing cumulative against the middle of the width class. From the resulting curve, Long first caught (Lc50) which is defined as the average total length where 50% of the fish are caught, taken according to the cumulative probability at 50%. This procedure follows the method by **Beverton dan Holt (1956**) dan **Sparre dan Venema (1998)**, as follows :
Lc50 = [TL – K(TL∞ – TL‟)]/ Z
Where , Lc50 is the first length, and T times caught where L 'is the average length of fish in the catch sample. Moreover, length at 75 and 95% fishing correlates with cumulative probability at 75 and 95%, respectively, as shown by **Pauly (1983).**
The result per recruit (Y / R) is determined according to the formula developed by **Beverton dan Holt (1957**), dan hasil relatif per rekrut (Y / R0) dihitung menurut **Pauly dan Soriano (1986).**

Data analysis

The FiSAT II tool was used to assess the population parameters, such asymptotic length (L∞), growth rate (K), natural mortality rate (M), fishing mortality rate (F), exploitation ratio (E), and relative yield per recruit indicators of samples from the assessed fish species encountered during the study period (**Gayanilo *et al.* 2005)**

RESULTS

Carapace width distribution

Carapace width distribution of blue swimming crab carapace (*P. pelagicus*) during the study from March 2023-August 2023, Who was caught with the bubu naga in Bawasalo waters Pangkep Regency to 1315 individuals on the group width carapace 6,70 cm – 13,80 cm, with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) (**Fig. 2**).




Fig (2). Carapace width distribution of blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

Carapace Width-weight relationship

The carapace width and weight relationship for males, females and combined sexes of *P. pelagicus* was described by equation as : $W= 0.1052L^{2.7245}$ (R2 = 7478), **(Fig 3**).




Fig (3). Width-weight Relationship of blue swimming crab (*P. pelagicus*) in Bawasalo waters

Age Group

Cohort analysis with Battacahrya's method *P. Pelagicus Consists of 3 age groups* is shown in **Fig (4).** Three cohort group were identified at length 8.66 sd 0.67, 10.27 sd 0.93, 12.16 sd 0.21 cm




Fig (4). Cohort groups by Bhattacharya's methode in Bawasalo Waters

Growth parameters

Estimated values of growth parameters were analyzed using the FISAT II program, The results of the analysis of the growth parameters estimation with the formula von Bertalannffy of blue swimming crab *P. pelagicus* including maximum length (L∞), growth coefficient (K) and age at birth (t0) that is L∞ = 15.1 cm, K = 0.41 per month and t0 = -0.48 **(Fig 5).**




Fig. (5). Growth curve of Von Bertalanffy blue swimming crab (*P. pelagicus*) in Bawasalo waters

Moralities and exploitation ratio

Mortality analysis based length-converted catch curve (**Fig. 6**) obtained total mortality rate, natural mortality, fishing mortality dan exploitation ratio blue swimming crab (*P. pelagicus*).




Fig. (6). Growth curve of Von Bertalanffy blue swimming crabs (*P. pelagicus*) in Bawasalo Waters

The total mortality (Z) was 2.70 yr^{-1}, natural mortality (M) and fishing mortality (F) was 1.22 yr^{-1} and 1.48 yr^{-1} and the exploitation ratio was 0.55 (**Table 1**).

Table 1. Value of population paramters blue swimming crab (*P. pelagicus*) in Bawasalo waters

Parameters	Value
Total mortality (Z)	2.70 yr-1
Natural mortality (M)	1.22 yr-1
Fishing mortality (F)	1.48 yr-1
Length at first capture (Lc)	8.56 cm
Length of recruitment (Lr)	8.22
Age at first capture (tc)	1.56
Age of recruitment (tr)	1.43
Exploitation ratio (E cur.)	0.55
The yield per recruit (Y/R)	20.6g
Biomass per recruit (B/R)	12.3g
The relative yield per recruit (Y/R')	0.009
Lc/L∞	0.567
M/K	2.976

Length at first capture

The length first capture (Lc50) is 10.92 cm. Length of catch probability 25% (Lc25) and 75% (Lc75) as 10.53 cm dan 11.22 cm **(Fig. 7).**




Fig. (7). Probability of capture

Yield per recruit (Y/R), and relative yield per recruit (Y'/R)

The yield per recruit (Y/R) for P. pelagicus was 20.6 gr pada current level mortality (F cur) 4.8 yr^{-1} (**Fig 8**). Relative yield per Y'/R recruit for different exploitation ratio (**Fig 9)**. From this curve, the maximum Y'/R is found at the exploitation ratio (Emax) = 0,50. The exploitation ratio E = 0.3.. would give the maximum Y/R' results in the protection of about 50% of the stock biomass




Fig. (8). Yield per recruit, biomass per recruit of blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig. (9). Relative yield per recruit and relative biomass per recruit in relation to different exploitation ratio of *P.pelagicus* in Bawasalo Waters

DISCUSSION

Carapace width distribution of blue swimming crab (*P. pelagicus*) during the study that is 6,70 cm – 13,80 cm, where the size of the carapace width < 10 cm as 66,69% and carapace width >10 cm as 33,31%**.** Research results of **Ihsan (2018)** in Pangkep's waters finding the dominant class interval males blue swimming crab on size 112-123 mm with a frequency of 265 heads (31,77%) and blue swimming crab females at intervals of classes 100-111 mm with a frequency of 223 heads (26,74%). Research results **Abdul Hamid (2019**) in Lasongko Bay shows carapace width on male blue swimming crab ranges from 4.1 cm -12.4 cm. Size of *P. pelagicus* obtained in Kwandang waters ranges from 7.0 – 17.5 cm (**Tirtadanu *et al.*, 2019**). Blue swimming crab catches obtained in Gresik and Lamongan dominated by the size class of carapace width of 11-12 cm and weight 70-90 g. The carapace width distribution shows that the carapace width capture >10 cm is 94%) **(Rahman *et al.*, 2019**). Crab populations in coastal waters of the northern Persian Gulf, Iran dominated by adult crabs (carapace width >100 mm) from December to May, juvenile crabs are dominant from June to October (**Safaie *et al.*,**

2015). The width of the crab carapace ranges from 60-150 mm (males) and 50-145 mm (females). The total weight males ranges to 48.0-275.50 g and females from 39.50 to 255.20 g **(Hosseini *et al.*, 2012)**. The length-weight relationship is one of the most important biological characters of fishes where weight of the fish increases as a function of its length (**Abeer *et al*, 2022**). **Fig (3)** exponent value (b) different to 3, this means the increase in carapace width is not as fast as the increase in weight. Coefficient b of 2.7344 shows growth allometrik negatif (P<0,05).

The same research was obtained **Muhsoni *et al* (2009)** in Bangkalan-Madura Waters, both males and females b<3. The same results were also obtained in the study **(Kanedi, 2020**) in the East Lampung Waters. The results are different from (**Ningrum, 2015**) which shows a value of b>3 (Allometrik positif) in the waters of Betahwalang and also at TPI Bulu, Jepara with growth pattern allometric positif **(Iksanti, 2022**). Carapace width-total weight relationship showed positive allometric growth (b-value= 3.10)(**El-Kasheif, 2021**). **Afzaal (2018)** reports the carapace length-weight of separate sexes (male and female) of *P. pelagicus* were also calculated as $W = 0.967L^{2.91}$ (R2 = 0.874) and carapace width-weight were as $W = 0.8138L^{2.95}$ (R2 = 0.882).

Blue swimming crab male and female crabs found in Bone waters consists of 1-3 cohorts (**Kembaren *et al.*, 2012**). The existence of cohort differences is suspected of subpopulation changes as a result of fishing pressure where fishing pressures of the sub population is decreasing (**Ernaningsih *et al.*, 2022**).

Value of the growth parameter Von Bertalanffy predicting the relationship between Fish stock estimation and fisheries resource management **(Abeer *et a*l, 2022).** The value of K indicates the growth rate blue swimming crab slow and takes a long time to reach its maximum length. Different growth coefficients in crustaceans can be due to several factors that is age, genetics, and seasons that affect water temperature (**Sara *et al.*, 2016**). Results of research on growth parameters of blue swimming crab (*P. pelagicus*) At different locations and times show different values (**Table. 2**).

Table 2. Growth parameter of blue swimming crabs (*P. pelagicus*) at different locations and times.

Researcher/year	Location	K (yr^{-1})	Lmaks (cm)
Kembaren/2012	Bone Bay	1,27	15.9
Sara/2017	Southeast Sulawesi Waters	0.55	16.9
Tirtadanu/2019	Indonesia	1,24	-
Abrenica/2021	Philipines	1,30	21,65
Rabaoui/2021	Saudi Arabian	2,12	-
Kasheif, 2021	Red Sea, Egypt	0,41	21.19
El-Betar/2022	Egypt	1,40	16.60
The present study/2023	Bawasalo, Pangkep	0,41	15.10

Total mortality rate (Z), natural mortality (M) dan fishing mortality (F) this study is lower than obtained by **Iksanti, *et al* (2022).** found a blue swimming crab landed at TPI Jepara at total mortality rate (Z) 2.99/year, natural mortality (M) 1,31/year dan fishing mortality (F) sebesar 1,68/year. Total mortality rate (Z), natural mortality (M) and fishing mortality (F) *of P. pelagicus* caught in Bardawil Wetland, Egypt 4.68/year, 2.25/year and 2.43/year respectively (**El-Betar, 2022**). The total mortality, natural mortality and capture mortality rates *of P. pelagicus* found in Egypt's Red Sea waters were 2,929/year, 1,285/year and 1,644/year, respectively (**El-Kasheif, 2021).** Fish species are more susceptible to activities fishing rather than deaths caused naturally by predation, competition, and disease (**Ofori-Danson *et al,* 2021**). The exploitation pressure of a fishing area can also be related to the parameters the dynamics of the fish population (**Ernaningsih *et al,* 2022**).

The ratio of exploitation is very important to estimate the state of the stock is optimal, under exploitation or over exploitation (**Mehanna *et al.,* 2019**). The ratio of exploitation of *P. pelagicus* found in this study in Bawasalo waters Pangkep Regency of 0.55 indicates that exploitation ratio exceeds the optimal limit (MSY). The results of interviews with local fishermen that the capture of blue swimming crab the intensity is quite high where fishermen catch every day using several types of fishing gear. In the Tiworo Strait shows that the population of blue swimming crab in the category of moderate exploitation (**Permatahati *et al,* 2020**).

CONCLUSION

Distribution carapace width 6.70 cm – 13.80 cm with the highest proportion at carapace width 9.10 – 9.60 cm (19,7%) and the lowest carapace width is 13.3 – 13.8 cm (0,2%.) where the carapace width capture < 10 cm with a proportion of 66.69% and carapace width >10 cm with a proportion of 33.31%. carapace width-weights relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric), the growth rate is low, the mortality rate of arrest is greater than the natural mortality rate and the exploitation ratio at limit of maksimal eksploitation (MSY) .

REFERENCES

Abrenica, B. T. (2021). Stock Assessment of the Blue Swimming Crab *Portunus pelagicus* (Linnaeus, 1758) in Stock Enhancement Sites of Danajon Bank, Central Philippines. *Philippine Journal of Fisheries*, *28*(2), : 210–227. https://doi.org/10.31398/tpjf/28.2.2020C0014

Afzaal, Z.; Kalhoro, M.; Buzdar, M.; Tariq, S., Shafi, M.; Nadeem, A.; Imran, S.; Saeed, F.; Sohail, M.; Hassan, R.; Haroon, A.; Shah, H. and Ahmed, I. (2018). Carapace length-weight and carapace width-weight relationship of *Portunus pelagicus* (Linnaeus, 1758) in Pakistani waters northern Arabian Sea. *Indian Journal of Geo-Marine Sciences*, *47*(4), 890–896.

https://api.elsevier.com/content/abstract/scopus_id/85045961734

Agus, S. B.; Zulbainarni, N.; Sunuddin, A.; Subarno, T.; Nugraha, A. H.; Rahimah, I.; Alamsyah, A.; Rachmi, R. and Jihad. 2016. Spatial Distribution of blue swimming crabs (*Portunus pelagicus*) in the Eastern Season in the waters of Lancang Island, Thousand Islands. Indonesian Journal of Agricultural Sciences 21(3): 209-218.

Asiedu, B.; Amponsah, S. K. K.; Ofori-Danson, P. K. and Nunoo, F. K. E. (2023). Assessment of Exploited Population Dynamics of African Butter Catfish (Schilbe mystus) in Stratum VII of Lake Volta, Ghana. *Egyptian Journal of Aquatic Biology and Fisheries*, *27*(6), 153–168. https://doi.org/10.21608/ejabf.2023.328024

Beverton, R.J.H. and Holt, S.J. (1956). A review of methods for estimation of mortality rates in exploited fish populations with special reference to sources of bias in catch sampling. Rapp-v Reun. Cons Intern. Mer., 140:67-83.

El-Betar, T. A. (2022). Population structure and growth aspects of blue swimming crab, Portunus pelagicus, in Bardawil Wetland, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *26*(4), 885–903. https://doi.org/10.21608/ejabf.2022.255480

Ernaningsih.; Muh Jamal.; Hasnidar. And Hadijah. (2022). Population parameters and exploitation rates of tiger grouper (*Epinephelus fuscoguttatus*) in the Spermonde Islands of South Sulawesi Jurnal Bawal Widyariset Perikanan Tangkap. 14 (1) April 2022: 1-9. e-ISSN: 2502-6410.

El-Kasheif, M. A. (2021). Fishery Biology and Population Structure of the Blue Swimmer Crab, Portunus pelagicus, from the Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *25*(6), 269–283. https://doi.org/10.21608/EJABF.2021.211506

Hamid, A.; Wurdiatno, Y.; Batu, D. T. F. L. and Riani, E. (2015). Fecundity and Maturity of Rajungan Gonads (Portunus pelagicus) Females Incubate Eggs in Lasongko Bay,. Southeast Sulawesi. Bawal. 7(1), 43-50.

Hosseini, M.; Vazirizade, A.; Parsa, Y. and Mansori, A. (2012). Sex ratio, size distribution and seasonal abundance of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) in Persian Gulf Coasts, Iran. *World Applied Sciences Journal*, *17*(7), 919–925. https://www.scopus.com/inward/record.uri?eid=2-s2.0-84861959364&partnerID=40&md5=be3ff1b6f61442673ff55db3ab710992

Ihsan.; Eko, S. W.; Sugeng, H. W. and John, H. (2014). A Study of Biological Potential and Sustainability of Swimming Crab Population in the Waters of Pangkep Regency South Sulawesi Province. International Journal of Sciences: Basic and Applied Research. 16(1): 351-363.

Ihsan. (2018). Distribusi Ukuran Dan Pola Musim Penangkapan Rajungan (*Portunus pelagicus*) Di Perairan Kabupaten Pangkep, jurnal marine Fisheries, Hal: 73-83

Iksanti, R. M.; Redjeki, S. and Taufiq-Spj, N. (2022). Biological Aspects of blue swimming crab (*Portunus pelagicus*) Linnaeus, 1758 (Malacostraca: *Portunidae*). Reviewed from Morphometry and Gonad Maturity Level at TPI Bulu, Jepara. Journal of Marine Research, 11(3), 495-505.

I. Maiyza, S.; F. Mehanna, S. and A. El-karyoney, I. (2020). An evaluation for the exploitation level of Egyptian Marine Fisheries. *Egyptian Journal of Aquatic Biology and Fisheries*, *24*(7), 441–452. https://doi.org/10.21608/ejabf.2020.121292

Kanedi, M. M.; Rahardjo, P. and Maulita, M. (2020). Biological Aspects of blue

swimming crab (*Portunus pelagicus*) in Coastal East Lampung Regency,Provinsi Lampung. Buletin Jalanidhitah Sarva Jivitam, 2(1), 49-56.

Kembaren, D.D.; Ernawati, T. and Suprapto. (2012). Biology and population parameters of crabs (*Portunus pelagicus*) in Bone Waters and surrounding areas. Jurnal Balai Penelitian Laut Jakarta. Vol 18 No. 4. Desember 2012 : 273-281 (In Indonesia)

Kusuma, B. T. dan Dewi. D. M. (2017). The Role of Fisheries Policies and Institutions in Management of Knitting (P*ortunus pelagicus*) So that it becomes a sustainable fishery Prosiding Simposium Nasional Krustasea 2017.

Lin, Y. J.; Grandcourt, E. M.; Rabaoui, L.; Maneja, R. H.; Qurban, M. A.; Al-Abdulkader, K. and Roa-Ureta, R. H. (2019). Comparative analysis of life history traits and trends of abundance in coral reefs of the orange-spotted grouper (Epinephelus coioides) from two regions of the Arabian Gulf. *ICES Journal of Marine Science*, *76*(4), 987–998. https://doi.org/10.1093/icesjms/fsz006

Mehanna, S. F.; Makkey, A. F. and Desouky, M. G. (2018). Growth, mortality and relative yield per recruit of the sharptooth catfish clarias gariepinus (Clariidae) in Lake Manzalah, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *22*(5), 65–72.

Mehanna, S. F.; Osman, Y. A. A.; Khalil, M. T. and Hassan, A. (2019). Age and growth, mortality and exploitation ratio of epinephelus summana (Forsskål, 1775) and cephalopholis argus (schneider, 1801) from the egyptian red sea coast, hurghada fishing area. *Egyptian Journal of Aquatic Biology and Fisheries*, *23*(4), 65–75.

Nurdin, M.; S, Ali.; S. A, dan Yanuarita. D. (2015). Mortalitas dan Laju Eksploitasi Rajungan (*Portunus pelagicus*) di Perairan Pulau Salemo Kabupaten Pangkajene Kepulauan. Jurnal Ipteks PSP 2:316-321.

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016a). Sex Ratio and Size at First Maturity Of Blue Swimming Crab (Portunus pelagicus) at Salemo Island, South Sulawesi.Ilmu Kelautan 21:17-22

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016b).. Size Distribution and Growth Pattern of Rajungan (*Portunus pelagicus*) on Salemo Island, Pangkajene Islands Regency. at the National Symposium on Marine Affairs and Fisheries III Makassar 7 Mei 2016. Page : 315-322

Ningrum, V. P.; Ghofar, A. and Ain, C. (2015). Biological Aspects Of Blue Swimmer Crab (*Portunus Pelagicus*) In Betahwalang Waters And Around. Saintek Perikanan: Indonesian Journal Of Fisheries Science And Technology, 11(1), 62-71.

Ofori-Danson, P.K.; Asiedu, B.; Alhassan, E.H.; Atsu, D.K. and Amponsah, S.K.K. (2021). Pre-Impoundment Fish Stock Assessment of the Black Volta: A Contribution to Fisheries Management of Bui Reservoir in Ghana. West African Journal of Applied Ecology, vol. 29(1): 35 – 48.

Pauly, D. (1979). Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries (No. 1; ICLARM, ICLARM Studies and Reviews). https://digitalarchive.worldfishcenter.org/ handle/ 20.500.12348/3693.

Pauly, D. (1980). On the interrelationship between natural mortality, growth

parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Intern.Explor. Mer., 39(2): 175-192.

Pauly, D. (1983). Some simple methods for the assessment of tropical fish stocks. FAO, Fish. Tech. Pap., 234: pp. http://www.fao.org/DOCREP/003/X6845E/X684 5E00.HTM

Rahman, M. A.; Iranawati, F.; Sambah, A. B. and Wiadnya, D. G. R. (2019). Biological condition and carapace width frequency distribution of blue swimming crabs (Portunus pelagicus) in Gresik and Lamongan, East Java. *IOP Conference Series: Earth and Environmental Science*, *370*(1). https://doi.org/10.1088/1755-1315/370/1/012010

Permatahati, Y. I.; Bugis, N. N.; Sara, L. and Hasuba, T. F. (2020). Stock status of blue swimming crab (Portunus pelagicus Linnaeus, 1758) in Tiworo Strait waters, Southeast Sulawesi, Indonesia. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, *25*(2), 85–90. https://doi.org/10.14710/ik.ijms.25.2.85-90

Rabaoui, L. (2021). Distribution, abundance, and life history traits of the blue swimming crab Portunus segnis (Forskål, 1775) in the Saudi waters of the Arabian Gulf. *Regional Studies in Marine Science*, *46*. https://doi.org/10.1016/j.rsma.2021.101895

Russ, G.R. 1991. Coral Reef Fisheries: Effect and Yields. In:The Ecology of Fishes On Coral Reef. Ed. P.F Sale, pp.601-35. Academic Press Limited, Lonndon.

Safaie, M.; Shokri, M. R.; Kiabi, B. H. and Pazooki, J. (2015). Biomass, CPUE and size frequency distribution of blue swimming crab Portunus segnis (Forskal, 1775) in coastal waters of the northern Persian Gulf, Iran. *Journal of the Marine Biological Association of the United Kingdom*, *95*(4), 763–771. https://doi.org/10.1017/S0025315414001635

Sara, L. and Astuti, O. (2019). Status of blue swimming crab (Portunus pelagicus) population in southeast sulawesi waters, Indonesia. *AACL Bioflux*, *12*(5), 1909–1917. https://www.scopus.com/inward/record.uri?eid=2-s2.0-85075242107&partnerID=40&md5=b0dc93ae0c32e8fa7812b1e9b52d2e3a

Sparre, P. and Venema, S. C. (2005). Introduction to TropicalFish Stock Assesment (Part 1, p. 376)

Sujatha, K.; Shrikanya, K. V. L. and Iswarya Deepti, V.A. (2015). Species diversity and some aspects ofreproductive biology and life history of groupers(Pisces: Serranidae: Epinephelinae) off the centraleastern coast of India. Marine Biology Research,11(1), 18–33. https://doi.org/10.1080/17451000.2014.949271

Tirtadanu. (2019). Fishery, population parameters and exploitation status of blue swimming crab (Portunus pelagicus) in kwandang waters, indonesia. *AACL Bioflux*, *12*(4), 1323–1334. https://api.elsevier.com/content/abstract/scopus_id/85073378715

Then, A. Y.; Hoenig, J. M. and Huynh, Q. C. (2018). Estimating fishing and natural mortality rates, andcatchability coefficient, from a series of observationson mean length and fishing effort. ICES Journal ofMarine Science, 75(2), 610–620. https://doi.org/10.1093/icesjms/fsx177.

Wiyono, E. S. (2015). The dynamic of landing blue swimming crab (Portunus pelagicus) catches in Pangkajene Kepulauan, South Sulawesi, Indonesia. *AACL Bioflux*, *8*(2), 134–141. https://api.elsevier.com/content/abstract/scopus_id/84925943854

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

1109: 4089-P

2 messages

Magdy Khalil <mtkhalil52@hotmail.com> Sat, Apr 27, 2024 at 11:46 PM
To: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Dear Authors,
Your article is in the final step; English language reviewing.
Regards

From: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>
Sent: Saturday, April 27, 2024 11:45 AM
To: Magdy Khalil <mtkhalil52@hotmail.com>
Subject: Re: 4089-P

dear
I sent a citation correction
Yours faithfully
Ernaningsih

On Sat, Apr 27, 2024 at 2:41 PM Magdy Khalil <mtkhalil52@hotmail.com> wrote:

> # Again....Do not waste our time
>
> **Dear Authors,**
> **Please revise carefully the attached version of your article before Language editing and publishing, especially the references, to match that of the Journal**;
>
> **Example**
> **Van der Geer, J.; Hanraads, J.A.J. ; Khalil, M. and Lupton, R.A.** (2010). The art of writing a scientific article. J. Sci. Commun., 163**:** 51–59.
>
> **Correct the chronological order of the references inside the text**;

From: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>
Sent: Saturday, April 27, 2024 7:03 AM
To: Magdy Khalil <mtkhalil52@hotmail.com>
Subject: Re: 4089-P

Dear Chief Editor
I am hereby sending back the journal which I have corrected according to his instructions. (Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
Yours faithfully
Ernaningsih

On Fri, Apr 26, 2024 at 5:34 PM Magdy Khalil <mtkhalil52@hotmail.com> wrote:

> **Dear Authors,**
> **Please revise carefully the attached version of your article before Language editing and publishing, especially the references, to match that of the Journal**;
>
> **Example**

Van der Geer, J.; Hanraads, J.A.J. ; Khalil, M. and Lupton, R.A. (2010). The art of writing a scientific article. J. Sci. Commun., 163**:** 51–59.

et al. should be written italic
Line space of the text is 1.15
Correct the chronological order of the references inside the text;

Correct Fig. 7 and organize the whole article

From: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>
Sent: Wednesday, April 17, 2024 3:16 AM
To: Magdy Khalil <mtkhalil52@hotmail.com>
Subject: Re: 4089

Dear chief editor
I sent the revised journal as directed (title: Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
and I send proof of journal payment (ID EJABF-2403-4089)
Yours faithfully
Ernaningsih

On Tue, Apr 16, 2024 at 6:38 AM Magdy Khalil <mtkhalil52@hotmail.com> wrote:

Dear Authors,
Please revise carefully the attached version of your article before Language editing and publishing, especially the references, to match that of the Journal;

Example
Van der Geer, J.; Hanraads, J.A.J. ; Khalil, M. and Lupton, R.A. (2010). The art of writing a scientific article. J. Sci. Commun., 163**:** 51–59.

Correct the chronological order of the references inside the text;

Please Send the cost of publication ($160) to Magdy Tawfik Khalil Ibrahim
Commercial International Bank (CIB)
Address: Thawra Branch- Cairo, Egypt
Account No. 100016289684
SWIFT CODE: CIBEEGCX081
IBAN: EG450010000600000100016289684
Beneficial address: Faculty of Science, Ain Shams University, Cairo, Egypt
Please write your full name in the receipt of the Bank

From: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>
Sent: Monday, April 15, 2024 12:36 PM
To: mtkhalil52@hotmail.com <mtkhalil52@hotmail.com>
Subject: Manuscript ID EJABF-2403-4089

dear Chief-editor
I am sending a journal manuscript which I have revised according to the Reviewer's suggestions with the title Population parameters and exploitation ratio of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) from the Bawasalo waters Pangkep Regency, South Sulawesi). (ID EJABF-2403-4089)
regards.
Ernaningsih

4089 P (1).doc
1103K

Ernaningsih Aras <ernaningsih.aras@umi.ac.id> Sat, May 18, 2024 at 7:01 PM

To: Magdy Khalil <mtkhalil52@hotmail.com>

Dear
Chef-Editor.
what is the position of my journal.
Thank You
Ernaningsih (1109: 4089-P)
[Quoted text hidden]

mail-UMI
e-mail Kampus UMI

Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Issue 16

1 message

Magdy Khalil <mtkhalil52@hotmail.com> Thu, May 23, 2024 at 6:13 PM
To: Ernaningsih Aras <ernaningsih.aras@umi.ac.id>

Dear Authors,
Congratulations….Your article has been published in the EJABF Journal.
Regards

Issue 16-4089 P (1) (1).pdf
1034K

Egyptian Journal of Aquatic Biology & Fisheries
Zoology Department, Faculty of Science,
Ain Shams University, Cairo, Egypt.
ISSN 1110 – 6131
Vol. 28(3): 285 – 298 (2024)
www.ejabf.journals.ekb.eg



Population Parameters and Exploitation Ratio of Blue Swimming Crab, *Portunus pelagicus* (Linnaeus, 1758) from the Bawasalo Waters Pangkep Regency, South Sulawesi

Ernaningsih*[1], Danial[1], Syahrul[1], Moh.Tauhid Umar[2], Nursyahran[3]

[1]Faculty of Fisheries and Marine Science Universitas Muslim Indonesia Makassar 90231
[2]Faculty of Marine Science and Fisheries, Hasanuddin University Makassar
[3]Institute of Maritime Technology and Business Balikdiwa, Makassar 90245
Corresponding Author: ernaningsih.aras@umi.ac.id

ARTICLE INFO

Article History:
Received: March 13, 2024
Accepted: April 23, 2024
Online: May 21, 2024

Keywords:
Portunus pelagicus,
Growth,
Mortalities,
Exploitation ratio,
Bawasalo waters

ABSTRACT

A total of 1315 individuals were collected from the catches off the Bawasalo waters during March -August 2023. This study aimed to analyze the population parameters and exploitation rate of blue swimming crab (*Portunus pelagicus*), caught by fishermen using a bubu naga fishing gear in the waters of Bawasalo, Pangkep District, South Sulawesi. The observed parameters included the size distribution of the carapace width, carapace width-weights relationship, age group, growth rate, total mortality rate, natural rate, fishing rate, and exploitation ratio analyzed with the ELEFAN II program. The size of carapace width ranged from 6.70 to 13.80cm, with the highest proportion observed at carapace widths between 9.10 and 9.60cm (19.7%) and the lowest proportion observed at carapace widths between 13.3 and 13.8cm (0.2%). Carapace widths below 10cm accounted for 66.69% of captures, while those above 10cm accounted for 33.31%. The carapace width-weight relationship was computed as $W = 0.1038L^{2.7334}$ (negative allometric). The von Bertalanffy growth parameters were estimated to be $K = 0.41\ yr^{-1}$, Lmax = 15.10 cm, and t0 = -0.723 yr. The total mortality (Z), natural mortality (M), and fishing mortality (F) were calculated as 2.70, 1.22, and 1.48 yr^{-1}, respectively. The exploitation rate (E = 0.55) suggests that the stock of *P. pelagicus* was slightly higher than that of an optimally exploited stock.

INTRODUCTION

Blue swimming crab is widely distributed throughout Indonesia **(Kusuma dan Dewi, 2017)**. In several regions of Indonesia, blue swimming crab is used as one of the people's fishery businesses including the Pangkep Regency **(Ihsan *et al*., 2014; Nurdin *et al*., 2016)**, Lasongko Bay **(Hamid *et al*., 2015)** and the Thousand Islands **(Agus *et al*., 2016)**.

Indexed in Scopus ELSEVIER DOAJ IUCAT

Pangkep Regency, located in south Sulawesi, has waters with considerable blue swimming crab (*P. pelagicus*) that has the potential to became one of the important commodities. In Pangkep Regency itself, one of the blue swimming crab producing centers is on Salemo Island **(Nurdin *et al.*, 2016b)**. The total exports of blue swimming crab in 2020 reached 27,616,332 tons, with an export value reaching USD 367,519,713 **(KKP, 2021)**. It ranks fourth in the total export value of Indonesian fishery products after shrimp, tuna and squid / cuttlefish / octopus.

The high export value indicates high production activities and exploitation of blue swimming crab, supported by open access fishing activities. Blue swimming crab (*P. pelagicus*) is currently under considerable pressure as the volume and size of the catch decreases **(Wiyono, 2015)**. High fishing pressure from blue swimming crab (*P. pelagicus*) in some areas in Indonesia has led to a reduction in stocks **(Tirtadanu, 2019)**. The catch and abundance of blue swimming crab on Salemo Island, Pangkep Regency, has decreased affecting the income of fishermen **(Nurdin *et al.*, 2015)**. Fisheries resources have seriously declined due to overfishing, illegal fishing activities, and pollution **(Mehanna *et al.*, 2018; Lin *et al.*, 2019; Asiedu *et al.*, 2023)**.

Biological information and population dynamics are radical for stock estimation **(Then *et al.*, 2018)**. Moreover, information regarding biological aspects is a major factor for management purposes **(Sujatha *et al.*, 2015)**. Therefore, for management purposes, information on fisheries biology and population structure is required **(El-Kasheif, 2021)**. The sustainability of the blue swimming crab requires population biology information and estimating fishery stocks as a scientific basis study for the availability of sustainable resources **(Froese *et al*, 2011; Maiyza *et al.*, 2020**). Research related to blue swimming crab in Pangkep Regency has been made regarding the arrest **(Ihsan, 2018)**. Bawasalo waters is one of the blue swimming crab fishing areas in Pangkep Regency. However, studies on the population parameters and exploitation ratio of blue swimming crab (*P. pelagicus*) caught with bubu naga in Bawasalo waters have not yet been conducted. Based on this, it was important to conduct research related to population parameters and exploitation rates which included the distribution of carapace width, carapace width-weight relationship, age group, growth rate, mortality rate, and rate of exploitation. These atudies are crucial for assessing the sustainability of blue swimming crab resources.

MATERIALS AND METHODS

A total of 1159 blue swimming crabs (*P. pelagicus*) were randomly drawn from March 2023 to August 2023 from the catch of bubu naga originating from Bawasalo waters, Segeri District, Pangkep Regency (Fig. 1). Each sample of blue swimming crab was measured for carapace width (CW) in cm (The width of the blue swimming crab is measured from the longest lateral spines on the sides of its body) and weight (W) in g.




Fig. 1. Map of research location of *P. pelagicus* in Bawasalo waters

Length frequency data were analyzed by using the Bhattacharya's method **(Bhattacharya, 1967)** that was involved in the FiSAT II software package **(Gayanilo, 2005)**.

The total length and body weight of fish were used for the length-weight relationship. The length-weight relationship was analyzed using the equation below **(Ricker, 1975)**:

$$W = aLb$$

Where, W = Total body weight of the fish (g); L = Total carapace width (cm); "a" = Regression constant, and "b" = Koefisien regresi; Value "b" provide information about the type of fish growth. The degree of association between length-weight variables was calculated by the correlation coefficient (R2), and the degree of its statistical significance was assessed following the method indicated by **Santos *et al*. (2002)**, as follows:

$$\text{Log } W = \text{Log } a + b \text{ Log } TL$$

If b = 3, the growth is isometric, b ≠ 3, the growth is allometric (allometric negative if b < 3, and the allometric is positive if b > 3).

Growth parameters followed the growth function of von Bertalanffy (VBGF) including growth rate (K) and asymptotic length (L∞) with equations.

$$L_t = L_\infty\left(1 - e^{-K(t - t_0)}\right)$$

Where, Lt is the average length at time (or age) t, L ∞ is the asymptotic mean length; K is the coefficient of growth rate (per year), and t; represents time or age. The theoretical age at zero length (t0) followed the equation shown by **Pauly (1979)**, as follows:

$$\text{Log } 10 (-t0) = -0.3922 - 0.2752 \log 10 L\infty - 1.038 \log 10 K$$

Estimation of total mortality (Z) was carried out by the length converted catch curve method **(Pauly, 1983)**. Natural mortality rate **(M) = Log M = -0.0066 – 0.279 log L∞ + 0.6543 log K + 0.4634 log T (Pauly, 1980)**, with a flat temperature of 30^0C. Fishing mortality rate (F)= Z-M. Exploitation rate was calculated using the equation described by **Gulland (1971)**: **E = F/Z**.

Long was allegedly first caught based on the cumulative probability of fishing effort across the middle of the width class. From the resulting curve, Long was first caught (Lc50), defined as the average total length where 50% of the fish are caught, taken according to the cumulative probability at 50%. This procedure follows the method described by **Beverton dan Holt (1956)** and **Sparre dan Venema (1998)**, as follows:

Lc50 = [TL – K(TL∞ – TL‟)]/ Z

Where, Lc50 is the first length, and T times caught, where L 'is the average length of fish in the catch sample. Moreover, length at 75 and 95% fishing correlates with cumulative probability at 75 and 95%, respectively, as shown by **Pauly (1983)**.

The result per recruit (Y/ R) was determined according to the formula developed by **Beverton dan Holt (1957**), and the relative per recruit (Y / R0) was calculated according to **Pauly and Soriano (1986)**.

Data analysis

The FiSAT II tool was used to assess the population parameters, such asymptotic length (L∞), growth rate (K), natural mortality rate (M), fishing mortality rate (F), exploitation ratio (E), and relative yield per recruit indicators of samples from the assessed fish species encountered during the study period **(Gayanilo *et al.*, 2005)**.

RESULTS

Carapace width distribution

During the study period from March 2023 to August 2023, a total of 1315 individuals of blue swimming crab (*P. pelagicus*) were caught with bubu naga in Bawasalo waters, Pangkep Regency. The carapace width distribution ranged from 6.70 to 13.80cm, with the highest proportion observed at carapace width, 9.10 to 9.60cm (19.7%), and the lowest proportion observed at carapace width, 13.3 to 13.8cm (0.2%) (Fig. 2).




Fig. 2. Carapace width distribution of blue swimming crabs (*P. pelagicus*) in Bawasalo waters

Carapace width-weight relationship

The carapace width and weight relationship for males, females and combined sexes of *P. pelagicus* was described in the following equation as : $W= 0.1052L^{2.7245}$ (R2 = 7478) (Fig. 3).




Fig. 3. Width-weight relationship of blue swimming crab (*P. pelagicus*) in Bawasalo waters

Age group

A cohort analysis using Battacahrya's method for *P. pelagicus* consists of three age groups, as shown in Fig. (4). Three cohort groups were identified with lengths of 8.66± 0.67, 10.27± 0.93, and 12.16± 0.21cm.




Fig. 4. Cohort groups using Bhattacharya's method in Bawasalo waters

Growth parameters

The estimated values of growth parameters were analyzed using the FISAT II program. The results of the analysis of the growth parameters estimation using the von

Bertalanffy formula for the blue swimming crab *P. pelagicus* included a maximum length (L∞) of 15.1cm, growth coefficient (K) of 0.41 per month, and age at birth (t0) of -0.48 (Fig. 5).




Fig. 5. Growth curve using von Bertalanffy blue swimming crab (*P. pelagicus*) in Bawasalo waters

Moralities and exploitation ratio

Mortality analysis based on the length-converted catch curve (Fig. 6) yielded the total mortality rate, natural mortality, fishing mortality, and exploitation ratio of the blue swimming crab (*P. pelagicus*).




Fig. 6. Growth curve using von Bertalanffy blue swimming crabs (*P. pelagicus*) in Bawasalo waters

The total mortality (Z) was 2.70 yr^{-1}, natural mortality (M) and fishing mortality (F) were 1.22 and 1.48 yr^{-1}, respectively, and the exploitation ratio was 0.55 (Table 1).

Table 1. Value of population paramters blue swimming crab (*P. pelagicus*) in Bawasalo waters

Parameters	Value
Total mortality (Z)	2.70 yr-1
Natural mortality (M)	1.22 yr-1
Fishing mortality (F)	1.48 yr-1
Length at first capture (Lc)	8.56 cm
Length of recruitment (Lr)	8.22
Age at first capture (tc)	1.56
Age of recruitment (tr)	1.43
Exploitation ratio (E cur.)	0.55
The yield per recruit (Y/R)	20.6g
Biomass per recruit (B/R)	12.3g
The relative yield per recruit (Y/R')	0.009
Lc/L∞	0.567
M/K	2.976

Length at first capture

The length at first capture (Lc50) was 10.92cm. The length of catch probability at 25% (Lc25) and 75% (Lc75) was 10.53 and 11.22cm, respectively (Fig. 7).




Fig. 7. Probability of capture

Yield per recruit (Y/R) and relative yield per recruit (Y'/R)

The yield per recruit (Y/R) for *P. pelagicus* was 20.6g at the current level of mortality (F cur) of 4.8 yr^{-1} (Fig 8). The relative yield per recruit (Y'/R) for different exploitation ratios is shown in Fig. (9). From this curve, the maximum Y'/R is found at the exploitation ratio (Emax) of 0.50. An exploitation ratio (E) of 0.3 would result in the maximum Y/R', protecting about 50% of the stock biomass.




Fig. 8. Yield per recruit, biomass per recruit of blue swimming crab (*P. pelagicus*) in Bawasalo waters




Fig. 9. Relative yield per recruit and relative biomass per recruit in relation to different exploitation ratio of *P. pelagicus* in Bawasalo waters

DISCUSSION

Carapace width distribution of blue swimming crab (*P. pelagicus*) during the study ranged between 6,70– 13,80cm. Approximately, 66.69% of the crabs had a carapace width of less than 10cm, while 33.31% had a carapace width greater than 10cm. In the research conducted by **Ihsan (2018)** in Pangkep waters, the dominant class interval for male blue swimming crabs was found to be 112- 123mm, with a frequency of 265 individuals (31.77%). For female blue swimming crabs, the dominant class interval was 100- 111mm, with a frequency of 223 individuals (26.74%). The research results of **Abdul Hamid (2019**) in Lasongko Bay show that the carapace width of male blue swimming crab ranges from 4.1- 12.4cm. The size of *P. pelagicus* obtained in Kwandang waters ranges from 7.0– 17.5cm **(Tirtadanu *et al.*, 2019)**. The catches of blue swimming crabs obtained in Gresik and Lamongan are dominated by individuals with a carapace width size class of 11- 12cm and a weight of 70- 90g. The carapace width distribution shows that the carapace width capture >10cm account for 94% **(Rahman *et al.*, 2019**). In the coastal waters of the northern Persian Gulf, Iran, crab populations are dominated by

adult crabs (carapace width >100mm) from December to May, while juvenile crabs are dominant from June to October **(Safaie *et al.*, 2015)**. The width of the crab carapace fluctuates from 60- 150mm (males) and 50- 145mm (females). The total weight of males ranges from 48.0- 275.50g, and for females from 39.50 to 255.20g **(Hosseini *et al.*, 2012)**. The length-weight relationship is one of the most important biological characters of fish, where weight of the fish increases as a function of its length **(Abeer *et al.*, 2022)**. In Fig. (3), the exponent value (b) is different from 3, indicating that the increase in carapace width is not as fast as the increase in weight. The coefficient b of 2.7344 indicates a negative allometric growth ($P< 0,05$).

The same research findings were obtained by **Muhsoni *et al.* (2009)** in Bangkalan-Madura waters; for both males and females b< 3. The same results were also obtained in the study by **Kanedi (2020)** in the East Lampung waters. However, these results differ from those of **Ningrum (2015)**, who showed a value of b> 3 (positive allometric) in the waters of Betahwalang and also at TPI Bulu, Jepara, indicating a positive allometric growth pattern **(Iksanti, 2022)**. Carapace width-total weight relationship showed a positive allometric growth (b-value= 3.10) **(El-Kasheif, 2021)**. **Afzaal (2018)** reported that, the carapace length-weight relationships for separate sexes (male and female) of *P. pelagicus* were calculated as $W = 0.967L^{2.91}$ ($R^2 = 0.874$), and the carapace width-weight relationships were calculated as $W = 0.8138L^{2.95}$ ($R^2 = 0.882$). Blue swimming crab male and female crabs found in Bone waters consists of 1- 3 cohorts (**Kembaren *et al.*, 2012**). The existence of cohort differences is suspected to be indicative of subpopulation changes resulting from fishing pressure, with the fishing pressure on the subpopulation decreasing (**Ernaningsih *et al.*, 2022**).

The value of the growth parameter von Bertalanffy predicts the relationship between fish stock estimation and fisheries resource management **(Abeer *et al.*, 2022)**. The value of K indicates that the growth rate of the blue swimming crab is slow, and it takes a long time to reach its maximum length. Different growth coefficients in crustaceans can be due to several factors that is age, genetics, and seasons that affect water temperature **(Sara *et al.*, 2016)**. The results of research on growth parameters of the blue swimming crab (*P. pelagicus*) at different locations and times exhibit varying values (Table 2).

Total mortality rate (Z), natural mortality (M) and fishing mortality (F) in this study are lower than those obtained by **Iksanti *et al.* (2022)**, who recorded a total mortality rate (Z) of 2.99/ year, natural mortality (M) of 1.31/ year, and fishing mortality (F) of 1.68/ year for blue swimming crabs that landed at TPI Jepara. The total mortality rate (Z), natural mortality (M) and fishing mortality (F) of *P. pelagicus* caught in Bardawil Wetland, Egypt, recorded 4.68/ year, 2.25/ year, and 2.43/ year, respectively **(El-Betar, 2022**). The total mortality, natural mortality and capture mortality rates of *P. pelagicus* found in Egypt's Red Sea waters were 2,929/ year, 1,285/ year and 1,644/ year, respectively **(El-Kasheif, 2021)**. Fish species are more susceptible to activities of fishing

rather than deaths caused naturally by predation, competition, and disease **(Ofori-Danson *et al.*, 2021)**. The exploitation pressure in a fishing area can also influence the dynamics of fish population parameters **(Ernaningsih *et al.*, 2022)**.

Table 2. Growth parameter of blue swimming crabs (*P. pelagicus*) in different locations and times

Researcher/year	Location	K (yr $^{-1}$)	Lmaks (cm)
Kembaren/2012	Bone Bay	1,27	15.9
Sara/2017	Southeast Sulawesi Waters	0.55	16.9
Tirtadanu/2019	Indonesia	1,24	-
Abrenica/2021	Philipines	1,30	21,65
Rabaoui/2021	Saudi Arabian	2,12	-
Kasheif, 2021	Red Sea, Egypt	0,41	21.19
El-Betar/2022	Egypt	1,40	16.60
The present study/2023	Bawasalo, Pangkep	0,41	15.10

The ratio of exploitation is very important to estimate the state of the stock is optimal, under exploitation or over exploitation **(Mehanna *et al.*, 2019)**. The ratio of exploitation of *P. pelagicus* found in this study in Bawasalo waters, Pangkep Regency, is 0.55, indicating that exploitation ratio exceeds the optimal limit (MSY). Interviews with local fishermen revealed that the intensity of blue swimming crab capture is quite high, with fishermen catching every day using several types of fishing gear. In the Tiworo Strait, the population of blue swimming crab is categorized as experiencing moderate exploitation **(Permatahati *et al.*, 2020)**.

CONCLUSION

The distribution of carapace width ranged from 6.70 to 13.80cm, with the highest proportion observed at carapace widths between 9.10 and 9.60cm (19.7%), and the lowest proportion at widths between 13.3 and 13.8cm (0.2%). Carapace widths less than 10cm accounted for 66.69% of the captures, while widths greater than 10cm accounted for 33.31%. The carapace width-to-weight relationship was computed as $W = 0.1038L^{2.7334}$, indicating a negative allometric growth. The growth rate is low, and the mortality rate of arrests exceeds the natural mortality rate. The exploitation ratio is at the limit of the maximum exploitation (MSY).

REFERENCES

Abrenica, B. T. (2021). Stock Assessment of the Blue Swimming Crab *Portunus pelagicus* (Linnaeus, 1758) in Stock Enhancement Sites of Danajon Bank, Central Philippines. *Philippine Journal of Fisheries*, *28*(2): 210–227. https://doi.org/10.31398/tpjf/28.2.2020C0014

Afzaal, Z.; Kalhoro, M.; Buzdar, M.; Tariq, S., Shafi, M.; Nadeem, A.; Imran, S.; Saeed, F.; Sohail, M.; Hassan, R.; Haroon, A.; Shah, H. and Ahmed, I. (2018). Carapace length-weight and carapace width-weight relationship of *Portunus pelagicus* (Linnaeus, 1758) in Pakistani waters northern Arabian Sea. *Indian Journal of Geo-Marine Sciences*, *47*(4): 890–896. https://api.elsevier.com/content/abstract/scopus_id/85045961734

Agus, S. B.; Zulbainarni, N.; Sunuddin, A.; Subarno, T.; Nugraha, A. H.; Rahimah, I.; Alamsyah, A.; Rachmi, R. and Jihad. 2016. Spatial Distribution of blue swimming crabs (*Portunus pelagicus*) in the Eastern Season in the waters of Lancang Island, Thousand Islands. Indonesian Journal of Agricultural Sciences 21(3): 209-218.

Asiedu, B.; Amponsah, S. K. K.; Ofori-Danson, P. K. and Nunoo, F. K. E. (2023). Assessment of Exploited Population Dynamics of African Butter Catfish (Schilbe mystus) in Stratum VII of Lake Volta, Ghana. *Egyptian Journal of Aquatic Biology and Fisheries*, *27*(6): 153–168. https://doi.org/10.21608/ejabf.2023.328024

Beverton, R.J.H. and Holt, S.J. (1956). A review of methods for estimation of mortality rates in exploited fish populations with special reference to sources of bias in catch sampling. Rapp-v Reun. Cons Intern. Mer., 140: 67-83.

El-Betar, T. A. (2022). Population structure and growth aspects of blue swimming crab, Portunus pelagicus, in Bardawil Wetland, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *26*(4): 885–903. https://doi.org/10.21608/ejabf.2022.255480

Ernaningsih.; Muh Jamal.; Hasnidar. And Hadijah. (2022). Population parameters and exploitation rates of tiger grouper (*Epinephelus fuscoguttatus*) in the Spermonde Islands of South Sulawesi Jurnal Bawal Widyariset Perikanan Tangkap. 14 (1) April 2022, 1-9. e-ISSN: 2502-6410.

El-Kasheif, M. A. (2021). Fishery Biology and Population Structure of the Blue Swimmer Crab, Portunus pelagicus, from the Red Sea, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *25*(6): 269–283. https://doi.org/10.21608/EJABF.2021.211506

Hamid, A.; Wurdiatno, Y.; Batu, D. T. F. L. and Riani, E. (2015). Fecundity and Maturity of Rajungan Gonads (Portunus pelagicus) Females Incubate Eggs in Lasongko Bay,. Southeast Sulawesi. Bawal. 7(1): 43-50.

Hosseini, M.; Vazirizade, A.; Parsa, Y. and Mansori, A. (2012). Sex ratio, size distribution and seasonal abundance of blue swimming crab, Portunus pelagicus (Linnaeus, 1758) in Persian Gulf Coasts, Iran. *World Applied Sciences Journal*, *17*(7): 919–925. https://www.scopus.com/inward/record.uri?eid=2-s2.0-84861959364&partnerID=40&md5=be3ff1b6f61442673ff55db3ab710992

Ihsan.; Eko, S. W.; Sugeng, H. W. and John, H. (2014). A Study of Biological Potential and Sustainability of Swimming Crab Population in the Waters of Pangkep Regency South Sulawesi Province. International Journal of Sciences: Basic and

Applied Research. 16(1): 351-363.

Ihsan. (2018). Distribusi Ukuran Dan Pola Musim Penangkapan Rajungan (*Portunus pelagicus*) Di Perairan Kabupaten Pangkep, jurnal marine Fisheries, Hal: 73-83

Iksanti, R. M.; Redjeki, S. and Taufiq-Spj, N. (2022). Biological Aspects of blue swimming crab (*Portunus pelagicus*) Linnaeus, 1758 (Malacostraca: *Portunidae*). Reviewed from Morphometry and Gonad Maturity Level at TPI Bulu, Jepara. Journal of Marine Research, 11(3): 495-505.

I. Maiyza, S.; F. Mehanna, S. and A. El-karyoney, I. (2020). An evaluation for the exploitation level of Egyptian Marine Fisheries. *Egyptian Journal of Aquatic Biology and Fisheries*, *24*(7): 441–452. https://doi.org/10.21608/ejabf.2020.121292

Kanedi, M. M.; Rahardjo, P. and Maulita, M. (2020). Biological Aspects of blue swimming crab (*Portunus pelagicus*) in Coastal East Lampung Regency,Provinsi Lampung. Buletin Jalanidhitah Sarva Jivitam, 2(1): 49-56.

Kembaren, D.D.; Ernawati, T. and Suprapto. (2012). Biology and population parameters of crabs (*Portunus pelagicus*) in Bone Waters and surrounding areas. Jurnal Balai Penelitian Laut Jakarta. 18 (4): 273-281 (In Indonesia)

Kusuma, B. T. dan Dewi. D. M. (2017). The Role of Fisheries Policies and Institutions in Management of Knitting (P*ortunus pelagicus*) So that it becomes a sustainable fishery Prosiding Simposium Nasional Krustasea 2017.

Lin, Y. J.; Grandcourt, E. M.; Rabaoui, L.; Maneja, R. H.; Qurban, M. A.; Al-Abdulkader, K. and Roa-Ureta, R. H. (2019). Comparative analysis of life history traits and trends of abundance in coral reefs of the orange-spotted grouper (Epinephelus coioides) from two regions of the Arabian Gulf. *ICES Journal of Marine Science*, *76*(4): 987–998. https://doi.org/10.1093/icesjms/fsz006

Mehanna, S. F.; Makkey, A. F. and Desouky, M. G. (2018). Growth, mortality and relative yield per recruit of the sharptooth catfish clarias gariepinus (Clariidae) in Lake Manzalah, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, *22*(5): 65–72.

Mehanna, S. F.; Osman, Y. A. A.; Khalil, M. T. and Hassan, A. (2019). Age and growth, mortality and exploitation ratio of epinephelus summana (Forsskål, 1775) and cephalopholis argus (schneider, 1801) from the egyptian red sea coast, hurghada fishing area. *Egyptian Journal of Aquatic Biology and Fisheries*, *23*(4): 65–75.

Nurdin, M.; S, Ali.; S. A, dan Yanuarita. D. (2015). Mortalitas dan Laju Eksploitasi Rajungan (*Portunus pelagicus*) di Perairan Pulau Salemo Kabupaten Pangkajene Kepulauan. Jurnal Ipteks PSP 2:316-321.

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016a). Sex Ratio and Size at First Maturity Of Blue Swimming Crab (Portunus pelagicus) at Salemo Island, South Sulawesi.Ilmu Kelautan 21, 17-22

Nurdin, M. S.; Ali, S. A. and Yanuarita. D. (2016b).. Size Distribution and Growth Pattern of Rajungan (*Portunus pelagicus*) on Salemo Island, Pangkajene Islands Regency. at the National Symposium on Marine Affairs and Fisheries III Makassar 7 Mei 2016. Page : 315-322

Ningrum, V. P.; Ghofar, A. and Ain, C. (2015). Biological Aspects Of Blue Swimmer Crab (*Portunus Pelagicus*) In Betahwalang Waters And Around. Saintek Perikanan:

Indonesian Journal Of Fisheries Science And Technology, 11(1): 62-71.

Ofori-Danson, P.K.; Asiedu, B.; Alhassan, E.H.; Atsu, D.K. and Amponsah, S.K.K. (2021). Pre-Impoundment Fish Stock Assessment of the Black Volta: A Contribution to Fisheries Management of Bui Reservoir in Ghana. West African Journal of Applied Ecology, vol. 29(1): 35 – 48.

Pauly, D. (1979). Theory and management of tropical multispecies stocks: a review, with emphasis on the Southeast Asian demersal fisheries (No. 1; ICLARM, ICLARM Studies and Reviews). https://digitalarchive.worldfishcenter.org/ handle/ 20.500.12348/3693.

Pauly, D. (1980). On the interrelationship between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Intern.Explor. Mer., 39(2): 175-192.

Pauly, D. (1983). Some simple methods for the assessment of tropical fish stocks. FAO, Fish. Tech. Pap., 234: pp. http://www.fao.org/DOCREP/003/X6845E/ X684 5E00.HTM

Rahman, M. A.; Iranawati, F.; Sambah, A. B. and Wiadnya, D. G. R. (2019). Biological condition and carapace width frequency distribution of blue swimming crabs (Portunus pelagicus) in Gresik and Lamongan, East Java. *IOP Conference Series: Earth and Environmental Science*, *370*(1). https://doi.org/10.1088/1755-1315/370/1/012010

Permatahati, Y. I.; Bugis, N. N.; Sara, L. and Hasuba, T. F. (2020). Stock status of blue swimming crab (Portunus pelagicus Linnaeus, 1758) in Tiworo Strait waters, Southeast Sulawesi, Indonesia. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, *25*(2): 85–90. https://doi.org/10.14710/ik.ijms.25.2.85-90

Rabaoui, L. (2021). Distribution, abundance, and life history traits of the blue swimming crab Portunus segnis (Forskål, 1775) in the Saudi waters of the Arabian Gulf. *Regional Studies in Marine Science*, *46*. https://doi.org/10.1016/j.rsma.2021.101895

Russ, G.R. 1991. Coral Reef Fisheries: Effect and Yields. In:The Ecology of Fishes On Coral Reef. Ed. P.F Sale, pp.601-35. Academic Press Limited, Lonndon.

Safaie, M.; Shokri, M. R.; Kiabi, B. H. and Pazooki, J. (2015). Biomass, CPUE and size frequency distribution of blue swimming crab Portunus segnis (Forskal, 1775) in coastal waters of the northern Persian Gulf, Iran. *Journal of the Marine Biological Association of the United Kingdom*, *95*(4): 763–771. https://doi.org/10.1017/S0025315414001635

Sara, L. and Astuti, O. (2019). Status of blue swimming crab (Portunus pelagicus) population in southeast sulawesi waters, Indonesia. *AACL Bioflux*, *12*(5): 1909–1917. https://www.scopus.com/inward/record.uri?eid=2-s2.0-85075242107&partnerID=40&md5=b0dc93ae0c32e8fa7812b1e9b52d2e3a

Sparre, P. and Venema, S. C. (2005). Introduction to TropicalFish Stock Assesment (Part 1, p. 376)

Sujatha, K.; Shrikanya, K. V. L. and Iswarya Deepti, V.A. (2015). Species diversity and some aspects ofreproductive biology and life history of groupers(Pisces: Serranidae: Epinephelinae) off the centraleastern coast of India. Marine Biology

Research,11(1): 18–33. https://doi.org/10.1080/17451000.2014.949271

Tirtadanu. (2019). Fishery, population parameters and exploitation status of blue swimming crab (Portunus pelagicus) in kwandang waters, indonesia. *AACL Bioflux*, *12*(4): 1323–1334. https://api.elsevier.com/content/abstract/scopus_id/85073378715

Then, A. Y.; Hoenig, J. M. and Huynh, Q. C. (2018). Estimating fishing and natural mortality rates, andcatchability coefficient, from a series of observationson mean length and fishing effort. ICES Journal ofMarine Science, 75(2): 610–620. https://doi.org/10.1093/icesjms/fsx177.

Wiyono, E. S. (2015). The dynamic of landing blue swimming crab (Portunus pelagicus) catches in Pangkajene Kepulauan, South Sulawesi, Indonesia. *AACL Bioflux*, *8*(2): 134–141. https://api.elsevier.com/content/abstract/scopus_id/84925943854